

MABUCHI AND AUBIN-YAU FUNCTIONALS OVER COMPLEX SURFACES

YI LI

*This paper is dedicated to my colleague and friend Lin Chen
who has passed away in an accident.*

ABSTRACT. In this note we construct Mabuchi $\mathcal{L}_\omega^M$ functional and Aubin-Yau functionals $\mathcal{I}_\omega^{AY}, \mathcal{J}_\omega^{AY}$ on any compact complex surfaces, and establish a number of properties. Our construction coincides with the original one in the Kähler case.

CONTENTS

1. Introduction	1
2. Err_ω map on complex manifolds	3
2.1. Err_ω map on compact complex manifolds	3
2.2. Vanishing property of Err_ω map on compact complex surface	6
3. Mabuchi $\mathcal{L}_\omega^M$ functional on complex surfaces	7
3.1. Mabuchi $\mathcal{L}_\omega^M$ functional on compact Kähler manifolds	7
3.2. Mabuchi $\mathcal{L}_\omega^M$ functional on compact complex surfaces	7
4. Aubin-Yau functionals on compact complex surfaces	11
4.1. Aubin-Yau functionals on compact Kähler manifolds	11
4.2. Aubin-Yau functionals over compact complex surfaces	13
References	16

1. INTRODUCTION

Let (X, ω) be a compact Kähler manifold of the complex dimension n . It's known that the volume V_ω depends only on the Kähler class of ω , namely,

$$\int_X (\omega + \sqrt{-1}\partial\bar{\partial}\varphi)^n = \int_X \omega^n$$

for any real-valued smooth function φ with $\omega_\varphi := \omega + \sqrt{-1}\partial\bar{\partial}\varphi > 0$, because of closedness of ω .

If (X, g) is a compact Hermitian manifold of the complex dimension n , the same result does not hold in general. In the Section 2, we consider a function to describe such a phenomena, i.e., we define

$$\text{Err}_\omega(\varphi) := \int_X \omega^n - \int_X \omega_\varphi^n,$$

where ω is its associated real $(1, 1)$ -form. The main result ¹ is

Proposition 1.1. *Let (X, g) be a compact Hermitian manifold of the complex dimension n and ω its associated real $(1, 1)$ -form. If $\partial\bar{\partial}(\omega^k) = 0$ for $k = 1, 2$, then*

$$\int_X \omega_\varphi^n = \int_X \omega^n =: V_\omega$$

for any real-valued function $\varphi \in C^\infty(X)_\mathbb{R}$ with $\omega_\varphi := \omega + \sqrt{-1}\partial\bar{\partial}\varphi > 0$.

In Kähler geometry, energy functionals, such as Mabuchi K -energy functional [6], Aubin-Yau energy functionals [7], and Chen-Tian energy functionals [1], play an important role in studying Kähler-Einstein metrics and constant scalar curvatures. When I was in Yau's Seminar, I asked myself that can we define energy functionals on compact complex manifolds? This is one motivation to write down this note. Another motivation comes from a question in S.-T. Yau's survey [12], that find necessary and sufficient conditions for a complex manifold to admit a Kähler structure. When $n = 2$, it was settled by Siu [8] or see [5]: A compact complex surface is Kähler if and only if its first Betti number is even. In the second part of this note we construct Mabuchi $\mathcal{L}_\omega^M$ functional and Aubin-Yau functionals $\mathcal{I}_\omega^{AY}, \mathcal{J}_\omega^{AY}$ on any compact complex surface.

Let (X, g) be a compact Hermitian manifold of the complex dimension 2 and ω be its associated real $(1, 1)$ -form. Let

$$\mathcal{P}_\omega := \{\varphi \in C^\infty(X)_\mathbb{R} | \omega_\varphi := \omega + \sqrt{-1}\partial\bar{\partial}\varphi > 0\}.$$

For any $\varphi', \varphi'' \in \mathcal{P}_\omega$, we define

$$\begin{aligned} \mathcal{L}_\omega^M(\varphi', \varphi'') &:= \frac{1}{V_\omega} \int_0^1 \int_X \dot{\varphi}_t \cdot \omega_{\varphi_t}^2 dt \\ &\quad - \frac{1}{V_\omega} \int_0^1 \int_X \sqrt{-1}\partial\omega \wedge (\bar{\partial}\dot{\varphi}_t \cdot \varphi_t) dt \\ &\quad + \frac{1}{V_\omega} \int_0^1 \int_X \sqrt{-1}\bar{\partial}\omega \wedge (\partial\dot{\varphi}_t \cdot \varphi_t) dt \end{aligned}$$

where φ_t is any smooth path in $\mathcal{P}_\omega$ from φ' to φ'' . Also we set

$$\mathcal{L}_\omega^M(\varphi) := \mathcal{L}_\omega^M(0, \varphi).$$

Theorem 1.2. *The functional $\mathcal{L}_\omega^M(\varphi', \varphi'')$ is independent of the choice of the smooth path $\{\varphi_t\}_{0 \leq t \leq 1}$ and satisfies the 1-cocycle condition. In particular*

$$\begin{aligned} \mathcal{L}_\omega^M(\varphi) &= \frac{1}{3V_\omega} \int_X \varphi(\omega^2 + \omega \wedge \omega_\varphi + \omega_\varphi^2) \\ &\quad + \frac{1}{2V_\omega} \int_X \varphi[-\sqrt{-1}\partial\omega \wedge \bar{\partial}\varphi + \sqrt{-1}\bar{\partial}\omega \wedge \partial\varphi]. \end{aligned}$$

Moreover, for any $\varphi \in \mathcal{P}_\omega$ and any constant $C \in \mathbb{R}$, we have

$$\mathcal{L}_\omega^M(\varphi, \varphi + C) = C \cdot \left(1 - \frac{\text{Err}_\omega(\varphi)}{V_\omega}\right);$$

for any $\varphi_1, \varphi_2 \in \mathcal{P}_\omega$ and any constant $C \in \mathbb{R}$, we have

$$\mathcal{L}_\omega^M(\varphi_1, \varphi_2 + C) = \mathcal{L}_\omega^M(\varphi_1, \varphi_2) + C \cdot \left(1 - \frac{\text{Err}_\omega(\varphi)}{V_\omega}\right).$$

¹V. Tosatti told the author that the same result has implicitly contained in [10] and [2].

In Section 4, we construct Aubin-Yau functionals on compact complex surfaces. Let (X, g) be a compact Hermitian manifold of complex dimension 2 and ω be its associated real $(1, 1)$ -form. Set

$$\begin{aligned}\mathcal{A}_\omega(\varphi) &:= \frac{1}{2V_\omega} \int_X \varphi \cdot -\sqrt{-1} \partial \omega \wedge \bar{\partial} \varphi, \\ \mathcal{B}_\omega(\varphi) &:= \frac{1}{2V_\omega} \int_X \varphi \cdot \sqrt{-1} \bar{\partial} \omega \wedge \partial \varphi.\end{aligned}$$

We define

$$\begin{aligned}\mathcal{I}_\omega^{\text{AY}}(\varphi) &:= \frac{1}{V_\omega} \int_X \varphi (\omega^2 - \omega_\varphi^2) + 2\mathcal{A}_\omega(\varphi) + 2\mathcal{B}_\omega(\varphi), \\ \mathcal{J}_\omega^{\text{AY}}(\varphi) &:= \frac{1}{V_\omega} \int_0^1 \int_X \varphi (\omega^2 - \omega_{s \cdot \varphi}^2) ds + \mathcal{A}_\omega(\varphi) + \mathcal{B}_\omega(\varphi).\end{aligned}$$

Theorem 1.3. *For any compact Hermitian manifold (X, g) of the complex dimension 2, we have*

$$\frac{1}{3} \mathcal{I}_\omega^{\text{AY}}(\varphi) \leq \mathcal{J}_\omega^{\text{AY}}(\varphi) \leq \frac{2}{3} \mathcal{I}_\omega^{\text{AY}}(\varphi)$$

for any $\varphi \in \mathcal{P}_\omega$, where ω is its associated real $(1, 1)$ -form.

We hope this exposition will give some ideas to study Yau's problem. The author is concerning the construction of those functionals on higher dimensional compact complex manifolds.

Acknowledgements. The author would like to thank Valentino Tosatti who read this note and pointed out a serious mistake in the first version. Furthermore, I also thank Chen-Yu Chi and Ming-Tao Chuan for useful discussion.

2. Err $_\omega$ MAP ON COMPLEX MANIFOLDS

Let (X, g) be a compact Hermitian manifold of the complex dimension n and write the associated real $(1, 1)$ -form as

$$\omega = \sqrt{-1} \cdot g_{i\bar{j}} \cdot dz^i \wedge d\bar{z}^j.$$

Let $\mathcal{P}_\omega$ be the space of all real-valued smooth functions $\varphi \in C^\infty(X)_\mathbb{R}$ so that $\omega + \sqrt{-1} \partial \bar{\partial} \varphi$ is positive definite on X :

$$\mathcal{P}_\omega := \{\varphi \in C^\infty(X)_\mathbb{R} \mid \omega + \sqrt{-1} \partial \bar{\partial} \varphi > 0\}.$$

Also we set

$$\mathcal{P}_\omega^0 := \left\{ \varphi \in \mathcal{P}_\omega \mid \sup_X \varphi = 0 \right\}.$$

2.1. Err $_\omega$ map on compact complex manifolds. To such a function $\varphi \in \mathcal{P}_\omega$ we associate the quantity

$$(2.1) \quad V_\omega(\varphi) := \int_X \omega_\varphi^n$$

the volume of X with respect to φ . In particular we set

$$(2.2) \quad V_\omega := V_\omega(0) = \int_X \omega^n.$$

For X being Kähler, we have $V_\omega = V_\omega(\varphi)$ for any $\varphi \in \mathcal{P}_\omega$. In the non-Kähler case, it's not in general true. Hence it is reasonable to define

$$(2.3) \quad \text{Err}_\omega(\varphi) := V_\omega - V_\omega(\varphi) = \int_X \omega^n - \int_X \omega_\varphi^n.$$

A natural question is when does $\text{Err}_\omega(\varphi)$ vanish for any/a $\varphi \in \mathcal{P}_\omega$? Clearly there exists a smooth real-valued function $\varphi_0 \equiv 0 \in \mathcal{P}_\omega$ such that

$$\omega_{\varphi_0}^n = \omega^n, \quad \sup_X \varphi_0 = 0,$$

hence

$$(2.4) \quad \text{Err}_\omega(\varphi_0) = \int_X \omega^n - \int_X \omega^n = 0.$$

This gives us some information about $\text{Err}_\omega(\varphi)$ and motivates us to consider

$$(2.5) \quad \text{SupErr}_\omega := \sup_{\varphi \in \mathcal{P}_\omega^0} (\text{Err}_\omega(\varphi)), \quad \text{InfErr}_\omega := \inf_{\varphi \in \mathcal{P}_\omega^0} (\text{Err}_\omega(\varphi)).$$

In any case, one has

$$(2.6) \quad \text{InfErr}_\omega \leq 0 \leq \text{SupErr}_\omega \leq \int_X \omega^n.$$

It's interesting to find some conditions to guarantee that the equalities hold. To study this behavior of Err_ω we consider the following several natural conditions on ω :

• **Condition 1.1:**

$$(2.7) \quad \sqrt{-1}\partial\omega \wedge \bar{\partial}\omega \quad \text{and} \quad \sqrt{-1}\partial\bar{\partial}\omega \quad \text{are non-negative}$$

• **Condition 1.2:**

$$(2.8) \quad \sqrt{-1}\partial\omega \wedge \bar{\partial}\omega \quad \text{and} \quad \sqrt{-1}\partial\bar{\partial}\omega \quad \text{are non-positive}$$

• **Condition 2:**

$$(2.9) \quad \partial\bar{\partial}(\omega^k) = 0, \quad k = 1, 2.$$

• **Condition 3:**

$$(2.10) \quad d(\omega^{n-1}) = 0.$$

• **Condition 4:**

$$(2.11) \quad \partial\bar{\partial}(\omega^{n-1}) = 0.$$

Remark 2.1. Condition 2 was appeared in [4] as a sufficient condition to solving the complex Monge-Ampère equation on Hermitian manifolds. The metric satisfying the third condition is called a balanced metric, which naturally appears in string theory (V. Tosatti and B. Wenkove [9] solved the complex Monge-Ampère equation on Hermitian manifolds with balanced metrics; later, they [10] dropped off the balanced condition.); When $n = 2$, this condition is indeed the Kähler condition. A metric satisfying Condition 4 is referred to be a Gauduchon metric, and a theorem of Gauduchon [3] shows that there exists a Gauduchon metric on every compact Hermitian manifold. Notice that Condition 3 implies condition 4, and Condition 2 is equivalent to $\partial\bar{\partial}\omega = 0 = \partial\omega \wedge \bar{\partial}\omega$. In particular, Condition 2 implies Condition 1.1 and Condition 1.2. In our case $n = 2$, Condition 2 is equivalent to Condition 4.

Remark 2.2. For any two forms α of degree $|\alpha|$ and β of degree $|\beta|$, we have

$$\alpha \wedge \beta = (-1)^{|\alpha| \cdot |\beta|} \beta \wedge \alpha.$$

Also we have

$$d(\alpha \wedge \beta) = d\alpha \wedge \beta + (-1)^{|\alpha|} \alpha \wedge d\beta;$$

according to types, one deduces

$$\begin{aligned} \partial(\alpha \wedge \beta) &= \partial\alpha \wedge \beta + (-1)^{|\alpha|} \alpha \wedge \partial\beta, \\ \bar{\partial}(\alpha \wedge \beta) &= \bar{\partial}\alpha \wedge \beta + (-1)^{|\alpha|} \alpha \wedge \bar{\partial}\beta. \end{aligned}$$

Moreover, if $|\alpha| + |\beta| = 2n - 1$, then

$$\begin{aligned} \int_X \alpha \wedge \partial\beta &= (-1)^{|\beta|} \int_X \partial\alpha \wedge \beta = -(-1)^{|\alpha|} \int_X \partial\alpha \wedge \beta, \\ \int_X \alpha \wedge \bar{\partial}\beta &= (-1)^{|\beta|} \int_X \bar{\partial}\alpha \wedge \beta = -(-1)^{|\alpha|} \int_X \bar{\partial}\alpha \wedge \beta. \end{aligned}$$

Those formulae are useful in our computations.

Theorem 2.3. (i) If ω satisfies Condition 1.1, then

$$(2.12) \quad \text{InfErr}_\omega = 0.$$

(ii) Correspondingly, if ω satisfies Condition 1.2, then

$$(2.13) \quad \text{SupErr}_\omega = 0.$$

(iii) In particular $\text{SupErr}_\omega = \text{InfErr}_\omega = 0$ provided that ω satisfies Condition 2.

Proof. (i) we knew that $\text{Err}_\omega(\varphi_0) = 0$ for some $\varphi_0 \in \mathcal{P}_\omega^0$. To achieve the argument, we need only to show that $\text{Err}_\omega(\varphi) \geq 0$ for each function $\varphi \in \mathcal{P}_\omega^0$. By definition we have

$$\begin{aligned} &\text{Err}_\omega(\varphi) \\ &= - \int_X \omega_\varphi^n + \int_X \omega^n = \int_X -\sqrt{-1} \partial \bar{\partial} \varphi \wedge \sum_{i=0}^{n-1} \omega_\varphi^i \wedge \omega^{n-1-i} \\ &= \sum_{i=0}^{n-1} \int_X \omega_\varphi^i \wedge \omega^{n-1-i} \wedge (-\sqrt{-1} \partial \bar{\partial} \varphi) = \sum_{i=0}^{n-1} \int_X \sqrt{-1} \partial (\omega_\varphi^i \wedge \omega^{n-1-i}) \wedge \bar{\partial} \varphi \\ &= \sum_{i=0}^{n-1} \int_X [i \cdot \omega_\varphi^{i-1} \wedge \partial \omega \wedge \omega^{n-1-i} + \omega_\varphi^i \wedge (n-1-i) \omega^{n-2-i} \wedge \partial \omega] \wedge \sqrt{-1} \bar{\partial} \varphi \\ &= \sum_{i=0}^{n-1} \int_X [i \cdot \omega_\varphi^{i-1} \wedge \omega^{n-1-i} + (n-1-i) \omega_\varphi^i \wedge \omega^{n-2-i}] \wedge \partial \omega \wedge \sqrt{-1} \bar{\partial} \varphi \\ &= \sum_{i=0}^{n-1} \int_X \sqrt{-1} [\varphi \cdot (i \cdot \omega_\varphi^{i-1} \wedge \omega^{n-1-i} + (n-1-i) \cdot \omega_\varphi^i \wedge \omega^{n-2-i}) \wedge \bar{\partial} \partial \omega \\ &\quad + \varphi \cdot \bar{\partial} (i \cdot \omega_\varphi^{i-1} \wedge \omega^{n-1-i} + (n-1-i) \omega_\varphi^i \wedge \omega^{n-2-i}) \wedge \partial \omega] \\ &= \sum_{i=0}^{n-1} (I_i + II_i) \end{aligned}$$

where

$$\begin{aligned} I_i &= \int_X \varphi \cdot [i \cdot \omega_\varphi^{i-1} \wedge \omega^{n-1-i} + (n-1-i)\omega_\varphi^i \wedge \omega^{n-2-i}] \wedge (-\sqrt{-1}\partial\bar{\partial}\omega), \\ II_i &= \int_X \varphi \cdot \sqrt{-1}\partial[i \cdot \omega_\varphi^{i-1} \wedge \omega^{n-1-i} + (n-1-i)\omega_\varphi^i \wedge \omega^{n-2-i}] \wedge \partial\omega. \end{aligned}$$

Since $\sqrt{-1}\partial\bar{\partial}\omega \geq 0$ and $\varphi \leq 0$ on X , the first term I_i is non-negative. Applying the integration by parts to II_i , we deduce

$$\begin{aligned} II_i &= \int_X \varphi [i(i-1)\omega_\varphi^{i-2} \wedge \bar{\partial}\omega \wedge \omega^{n-1-i} + i(n-1-i)\omega_\varphi^{i-1} \wedge \omega^{n-2-i} \wedge \bar{\partial}\omega \\ &\quad + i(n-1-i)\omega_\varphi^{i-1} \wedge \bar{\partial}\omega \wedge \omega^{n-2-i} \\ &\quad + (n-1-i)(n-2-i)\omega_\varphi^i \wedge \omega^{n-3-i} \wedge \bar{\partial}\omega] \wedge \sqrt{-1}\partial\omega \\ &= \int_X \varphi \cdot \omega_\varphi^{i-2} \wedge \omega^{n-3-i} \wedge [i(i-1)\omega^2 + 2i(n-1-i)\omega_\varphi \wedge \omega \\ &\quad + (n-1-i)(n-2-i)\omega_\varphi^2] \wedge (-\sqrt{-1}\partial\omega \wedge \bar{\partial}\omega). \end{aligned}$$

Since $\sqrt{-1}\partial\omega \wedge \bar{\partial}\omega$ is non-negative and φ is non-positive, it follows that $II_i \geq 0$. Thus $\text{Err}_\omega(\varphi) \geq 0$ for each $\varphi \in \mathcal{P}_\omega^0$ and therefore $\text{InfErr}_\omega = 0$.

(ii) If ω satisfies Condition 1.2, the above proceeding gives that $\text{Err}_\omega(\varphi) \leq 0$ for each $\varphi \in \mathcal{P}_\omega^0$, i.e., $\text{SupErr}_\omega \leq 0$. Hence $\text{SupErr}_\omega = 0$.

(iii) It's an immediate consequence of (i) and (ii). $\square$

Corollary 2.4. *If ω satisfies Condition 2, then $\text{Err}_\omega(\varphi) = 0$ for any $\varphi \in \mathcal{P}_\omega^0$. Equivalently, in this case, the number $V_\omega(\varphi) = \int_X \omega_\varphi^n$ does not depend on the choice of $\varphi \in \mathcal{P}_\omega$ and equals $V_\omega = \int_X \omega^n$.*

2.2. Vanishing property of Err_ω map on compact complex surface. Let (X, g) be a Hermitian manifold of the complex dimension n and let ω_g be its associated real $(1, 1)$ -form. We say that g is a Gauduchon metric if $\partial\bar{\partial}(\omega_g^{n-1}) = 0$.

We recall a theorem of Gauduchon [3] or see Remark 2.1.

Theorem 2.5. (Gauduchon, 1984) *If X is a compact complex manifold of the complex dimension n , then in the conformal class of every Hermitian metric g there exists a Gauduchon metric g_G , i.e., there is a positive function $\varphi \in C^\infty(X)_\mathbb{R}$ such that $g_G := \varphi \cdot g$ is Gauduchon. If X is connected and $n \geq 2$, then g_G is unique up to a positive constants.*

Using the existence of Gauduchon metric, we can prove the following

Theorem 2.6. *Let (X, g) be a compact complex surface with Hermitian metric g and let ω_G be its associated Gauduchon metric. Then*

$$(2.14) \quad \text{Err}_{\omega_G}(\varphi) = 0$$

for all $\varphi \in \mathcal{P}_{\omega_G}$.

Proof. In what follows, we omit the subscript G and write ω_G as ω ; since ω is a Gauduchon metric, it follows that $\partial\bar{\partial}\omega = 0$. In the case of $n = 2$, we have

$$\begin{aligned} \text{Err}_\omega(\varphi) &= \int_X \omega^2 - \int_X \omega_\varphi^2 = \int_X (\omega - \omega_\varphi)(\omega + \omega_\varphi) \\ &= \int_X (\omega + \omega_\varphi) \wedge -\sqrt{-1}\partial\bar{\partial}\varphi \\ &= \int_X -\sqrt{-1}\partial\bar{\partial}(\omega + \omega_\varphi) \cdot \varphi \\ &= \int_X -2\sqrt{-1}\partial\bar{\partial}\omega \cdot \varphi = 0. \end{aligned}$$

Hence the theorem follows. $\square$

3. MABUCHI $\mathcal{L}_\omega^M$ FUNCTIONAL ON COMPLEX SURFACES

3.1. Mabuchi $\mathcal{L}_\omega^M$ functional on compact Kähler manifolds. Suppose that (X, ω) is a compact Kähler manifold of the complex dimension n . For any pair $(\varphi', \varphi'') \in \mathcal{P}_\omega \times \mathcal{P}_\omega$ we define

$$\mathcal{L}_\omega^M : \mathcal{P}_\omega \times \mathcal{P}_\omega \longrightarrow \mathbb{R}$$

as follows:

$$(3.1) \quad \mathcal{L}_\omega^M(\varphi', \varphi'') := \frac{1}{V_\omega} \int_0^1 \int_X \dot{\varphi}_t \cdot \omega_{\varphi_t}^n \cdot dt$$

where $\{\varphi_t : 0 \leq t \leq 1\}$ is any smooth path in $\mathcal{P}_\omega$ such that $\varphi_0 = \varphi'$ and $\varphi_1 = \varphi''$. For any $\varphi \in \mathcal{P}_\omega$ we set

$$(3.2) \quad \mathcal{L}_\omega^M(\varphi) := \mathcal{L}_\omega^M(0, \varphi).$$

Mabuchi [6] showed that the functional (3.1) is well-defined, and hence we can explicitly write down $\mathcal{L}_\omega^M(\varphi)$.

In this section we extend Mabuchi $\mathcal{L}_\omega^M$ functional to any compact complex surface by adding two extra terms on the right hand side of (3.1).

3.2. Mabuchi $\mathcal{L}_\omega^M$ functional on compact complex surfaces. Suppose now that (X, g) is a compact complex surface and ω be its associated real $(1, 1)$ -form. Let $\varphi', \varphi'' \in \mathcal{P}_\omega$ and $\{\varphi_t\}_{0 \leq t \leq 1}$ be a smooth path in $\mathcal{P}_\omega$ from φ' to φ'' .

Let

$$(3.3) \quad \mathcal{L}_\omega^0(\varphi', \varphi'') := \frac{1}{V_\omega} \int_0^1 \int_X \dot{\varphi}_t \cdot \omega_{\varphi_t}^2 \cdot dt.$$

Set

$$(3.4) \quad \psi(s, t) := s \cdot \varphi_t, \quad 0 \leq s \leq 1, \quad 0 \leq t \leq 1.$$

Consider a 1-form on $[0, 1] \times [0, 1]$

$$(3.5) \quad \Psi^0 := \left(\int_X \frac{\partial \psi}{\partial s} \cdot \omega_\psi^2 \right) ds + \left(\int_X \frac{\partial \psi}{\partial t} \cdot \omega_\psi^2 \right) dt.$$

Taking differential on Ψ^0 , we have

$$d\Psi^0 = I^0 \cdot dt \wedge ds$$

where

$$(3.6) \quad I^0 = \int_X \frac{\partial}{\partial t} \left(\frac{\partial \psi}{\partial s} \cdot \omega_\psi^2 \right) - \int_X \frac{\partial}{\partial s} \left(\frac{\partial \psi}{\partial t} \cdot \omega_\psi^2 \right).$$

Directly computing shows

$$\begin{aligned} I^0 &= \int_X \left[\frac{\partial^2 \psi}{\partial t \partial s} \cdot \omega_\psi^2 + \frac{\partial \psi}{\partial s} \cdot 2 \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \right] \\ &\quad - \int_X \left[\frac{\partial^2 \psi}{\partial s \partial t} \cdot \omega_\psi^2 + \frac{\partial \psi}{\partial t} \cdot 2 \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \right] \\ &= \int_X 2 \frac{\partial \psi}{\partial s} \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) - \int_X 2 \frac{\partial \psi}{\partial t} \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right). \end{aligned}$$

In the following we deduce two slightly different formulae of I^0 . The first one is

$$\begin{aligned} I^0 &= \int_X 2 \frac{\partial \psi}{\partial s} \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) + \int_X 2 \frac{\partial \psi}{\partial t} \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \\ &= \int_X -2 \sqrt{-1} \partial \left(\frac{\partial \psi}{\partial s} \cdot \omega_\psi \right) \wedge \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \\ &\quad + \int_X -2 \sqrt{-1} \bar{\partial} \left(\frac{\partial \psi}{\partial t} \cdot \omega_\psi \right) \wedge \partial \left(\frac{\partial \psi}{\partial s} \right) \\ &= \int_X -2 \sqrt{-1} \left[\partial \left(\frac{\partial \psi}{\partial s} \right) \wedge \omega_\psi + \frac{\partial \psi}{\partial s} \cdot \partial \omega \right] \wedge \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \\ &\quad + \int_X -2 \sqrt{-1} \left[\bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \wedge \omega_\psi + \frac{\partial \psi}{\partial t} \cdot \bar{\partial} \omega \right] \wedge \partial \left(\frac{\partial \psi}{\partial s} \right) \\ &= \int_X -2 \sqrt{-1} \frac{\partial \psi}{\partial s} \cdot \partial \omega \wedge \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) + \int_X -2 \sqrt{-1} \frac{\partial \psi}{\partial t} \cdot \bar{\partial} \omega \wedge \partial \left(\frac{\partial \psi}{\partial s} \right) \\ &= \int_X 2 \sqrt{-1} \frac{\partial \psi}{\partial s} \cdot \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \wedge \partial \omega + \int_X 2 \sqrt{-1} \frac{\partial \psi}{\partial t} \cdot \partial \left(\frac{\partial \psi}{\partial s} \right) \wedge \bar{\partial} \omega. \end{aligned}$$

Similarly, we have

$$I^0 = \int_X -2 \sqrt{-1} \frac{\partial \psi}{\partial s} \cdot \partial \left(\frac{\partial \psi}{\partial t} \right) \wedge \bar{\partial} \omega + \int_X -2 \sqrt{-1} \frac{\partial \psi}{\partial t} \cdot \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \wedge \partial \omega.$$

Next, we define

$$(3.7) \quad \mathcal{L}_\omega^1(\varphi', \varphi'') = \frac{1}{V_\omega} \int_0^1 \int_X a_2 \cdot \partial \omega \wedge (\bar{\partial} \dot{\varphi}_t \cdot \varphi_t) dt,$$

$$(3.8) \quad \mathcal{L}_\omega^2(\varphi', \varphi'') = \frac{1}{V_\omega} \int_0^1 \int_X b_2 \cdot \bar{\partial} \omega \wedge (\partial \dot{\varphi}_t \cdot \varphi_t) dt.$$

Here we require $\overline{a_2} = b_2$, and a_2, b_2 are determined later. As before, consider

$$\begin{aligned} \Psi^1 &= \left[\int_X a_2 \partial \omega \wedge \left(\bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \cdot \psi \right) \right] ds + \left[\int_X a_2 \partial \omega \wedge \left(\bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \cdot \psi \right) \right] dt, \\ \Psi^2 &= \left[\int_X b_2 \bar{\partial} \omega \wedge \left(\partial \left(\frac{\partial \psi}{\partial s} \right) \cdot \psi \right) \right] ds + \left[\int_X b_2 \bar{\partial} \omega \wedge \left(\partial \left(\frac{\partial \psi}{\partial t} \right) \cdot \psi \right) \right] dt. \end{aligned}$$

Therefore

$$d\Psi^1 = I^1 \cdot dt \wedge ds$$

where

$$(3.9) \quad \begin{aligned} I^1 &= \int_X a_2 \frac{\partial}{\partial t} \left[\partial \omega \wedge \left(\bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \cdot \psi \right) \right] \\ &\quad - \int_X a_2 \frac{\partial}{\partial s} \left[\partial \omega \wedge \left(\bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \cdot \psi \right) \right]. \end{aligned}$$

Dividing I^1 by a_2 yields

$$\begin{aligned} \frac{I^1}{a_2} &= \int_X -\frac{\partial}{\partial t} \left[\left(\bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \cdot \psi \right) \wedge \partial \omega \right] + \int_X \frac{\partial}{\partial s} \left[\left(\bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \cdot \psi \right) \wedge \partial \omega \right] \\ &= \int_X - \left[\bar{\partial} \left(\frac{\partial^2 \psi}{\partial t \partial s} \right) \cdot \psi + \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \cdot \frac{\partial \psi}{\partial t} \right] \wedge \partial \omega \\ &\quad + \int_X \left[\bar{\partial} \left(\frac{\partial^2 \psi}{\partial s \partial t} \right) \cdot \psi + \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \cdot \frac{\partial \psi}{\partial s} \right] \wedge \partial \omega \\ &= \int_X -\frac{\partial \psi}{\partial t} \cdot \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \wedge \partial \omega + \int_X \frac{\partial \psi}{\partial s} \cdot \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \wedge \partial \omega. \end{aligned}$$

In the same way, one deduces

$$d\Psi^2 = I^2 \cdot dt \wedge ds,$$

and

$$\frac{I^2}{b_2} = \int_X -\frac{\partial \psi}{\partial t} \cdot \partial \left(\frac{\partial \psi}{\partial s} \right) \wedge \bar{\partial} \omega + \int_X \frac{\partial \psi}{\partial s} \cdot \partial \left(\frac{\partial \psi}{\partial t} \right) \wedge \bar{\partial} \omega.$$

Combining above formulas, we have

$$(3.10) \quad -\frac{I^1}{a_2} + \frac{I^2}{b_2} = -\frac{I^0}{\sqrt{-1}}.$$

Setting $a_2 = -\sqrt{-1}$ and $b_2 = \sqrt{-1}$, we get

$$I^0 + I^1 + I^2 = 0.$$

Thus

$$(3.11) \quad d\Psi = 0$$

where

$$\Psi := \Psi^0 + \Psi^1 + \Psi^2.$$

The following theorem is an immediate consequence of the above discussion.

Theorem 3.1. *Let (X, g) be a compact complex surface and ω be its associated real $(1, 1)$ -form. The functional*

$$(3.12) \quad \begin{aligned} \mathcal{L}_\omega^M(\varphi', \varphi'') &:= \frac{1}{V_\omega} \int_0^1 \int_X \dot{\varphi}_t \cdot \omega_{\varphi_t}^2 dt \\ &\quad - \frac{1}{V_\omega} \int_0^1 \int_X \sqrt{-1} \partial \omega \wedge (\bar{\partial} \dot{\varphi}_t \cdot \varphi_t) dt \\ &\quad + \frac{1}{V_\omega} \int_0^1 \int_X \sqrt{-1} \bar{\partial} \omega \wedge (\partial \dot{\varphi}_t \cdot \varphi_t) dt \end{aligned}$$

is independent of the choice of the smooth path $\{\varphi_t\}_{0 \leq t \leq 1}$. In particular,

$$(3.13) \quad \begin{aligned} \mathcal{L}_\omega^M(\varphi) &:= \mathcal{L}_\omega(0, \varphi) = \frac{1}{3V_\omega} \int_X \varphi(\omega^2 + \omega \wedge \omega_\varphi + \omega_\varphi^2) \\ &+ \frac{1}{2V_\omega} \int_X \varphi[-\sqrt{-1}\partial\omega \wedge \bar{\partial}\varphi + \sqrt{-1}\bar{\partial}\omega \wedge \partial\varphi]. \end{aligned}$$

Proof. Applying Stocks' theorem to the region $\Delta = \{(s, t) \in \mathbb{R}^2 : 0 \leq s, t \leq 1\}$ and using equation (3.9), we have

$$\begin{aligned} 0 &= \int_\Delta d\Psi = \int_{\partial\Delta} \Psi = \int_{\partial\Delta} (\Psi^0 + \Psi^1 + \Psi^2) \\ &= \int_0^1 \dot{\varphi}_t \cdot \omega_{\varphi_t}^2 dt - \int_0^1 \int_X \frac{\partial\psi}{\partial s} \cdot \omega_\psi^2 ds \Big|_{t=0}^{t=1} \\ &+ \int_0^1 \int_X -\sqrt{-1}\partial\omega \wedge (\bar{\partial}\dot{\varphi}_t \cdot \varphi_t) dt \\ &- \int_0^1 \int_X -\sqrt{-1}\partial\omega \wedge \left(\bar{\partial} \left(\frac{\partial\psi}{\partial s} \right) \cdot \psi \right) ds \Big|_{t=0}^{t=1} \\ &+ \int_0^1 \int_X \sqrt{-1}\bar{\partial}\omega \wedge (\partial\dot{\varphi}_t \cdot \varphi_t) dt \\ &- \int_0^1 \int_X \sqrt{-1}\bar{\partial}\omega \wedge \left(\partial \left(\frac{\partial\psi}{\partial s} \right) \cdot \psi \right) ds \Big|_{t=0}^{t=1}. \end{aligned}$$

Equivalently,

$$\begin{aligned} \mathcal{L}_\omega^M(\varphi', \varphi'') &= \int_0^1 \int_X \frac{\partial\psi}{\partial s} \cdot \omega_\psi^2 ds \Big|_{t=0}^{t=1} \\ &+ \int_0^1 \int_X -\sqrt{-1}\partial\omega \wedge \left(\bar{\partial} \left(\frac{\partial\psi}{\partial s} \right) \cdot \psi \right) ds \Big|_{t=0}^{t=1} \\ &+ \int_0^1 \int_X \sqrt{-1}\bar{\partial}\omega \wedge \left(\partial \left(\frac{\partial\psi}{\partial s} \right) \cdot \psi \right) ds \Big|_{t=0}^{t=1}. \end{aligned}$$

It turns out that $\mathcal{L}_\omega^M(\varphi', \varphi'')$ is well-defined. For the second argument, we can choose the smooth path $\varphi_t = t \cdot \varphi$, $0 \leq t \leq 1$. $\square$

Remark 3.2. When (X, g) is a compact Kähler surface, the functional (3.12) or (3.13) coincides with the original one.

Definition 3.3. Suppose that S is a non-empty set and A is an additive group. A mapping $\mathcal{N} : S \times S \rightarrow A$ is said to satisfy the **1-cocycle condition** if

- (i) $\mathcal{N}(\sigma_1, \sigma_2) + \mathcal{N}(\sigma_2, \sigma_1) = 0$;
- (ii) $\mathcal{N}(\sigma_1, \sigma_2) + \mathcal{N}(\sigma_2, \sigma_3) + \mathcal{N}(\sigma_3, \sigma_1) = 0$.

Corollary 3.4. The functional $\mathcal{L}_\omega^M$ satisfies the 1-cocycle condition.

Corollary 3.5. For any $\varphi \in \mathcal{P}_\omega$ and any constant $C \in \mathbb{R}$, we have

$$(3.14) \quad \mathcal{L}_\omega^M(\varphi, \varphi + C) = C \cdot \left(1 - \frac{\text{Err}_\omega(\varphi)}{V_\omega} \right).$$

In particular, if $\partial\bar{\partial}\omega = 0$, then $\mathcal{L}_\omega^M(\varphi, \varphi + C) = C$.

Proof. We choose the smooth path $\varphi_t = \varphi + t \cdot C$, $t \in [0, 1]$. So

$$\begin{aligned} \mathcal{L}_\omega^M(\varphi, \varphi + C) &= \frac{1}{V_\omega} \int_0^1 \int_X C \cdot \omega_{\varphi+C \cdot t}^2 dt = \frac{1}{V_\omega} \int_0^1 \int_X C \cdot \omega_\varphi^2 dt \\ &= \frac{1}{V_\omega} \int_X C \cdot \omega_\varphi^2 = \frac{C}{V_\omega} \cdot V_\omega(\varphi) = C \cdot \left(1 - \frac{\text{Err}_\omega(\varphi)}{V_\omega}\right). \end{aligned}$$

If furthermore $\partial\bar{\partial}\omega = 0$, then $\text{Err}_\omega(\varphi) = 0$ for any $\varphi \in \mathcal{P}_\omega$. $\square$

Corollary 3.6. *For any $\varphi_1, \varphi_2 \in \mathcal{P}_\omega$ and any constant $C \in \mathbb{R}$, we have*

$$(3.15) \quad \mathcal{L}_\omega^M(\varphi_1, \varphi_2 + C) = \mathcal{L}_\omega^M(\varphi_1, \varphi_2) + C \cdot \left(1 - \frac{\text{Err}_\omega(\varphi_2)}{V_\omega}\right).$$

Proof. From Corollary 3.4, one has

$$\mathcal{L}_\omega^M(\varphi_1, \varphi_2 + C) + \mathcal{L}_\omega^M(\varphi_2, \varphi_1) = \mathcal{L}_\omega^M(\varphi_2, \varphi_2 + C).$$

Then the conclusion follows from Corollary 3.5. $\square$

4. AUBIN-YAU FUNCTIONALS ON COMPACT COMPLEX SURFACES

In this section we extend Aubin-Yau functionals to compact complex surfaces, including Kähler surfaces, and deduce a number of basic properties of these functionals.

4.1. Aubin-Yau functionals on compact Kähler manifolds. Suppose that (X, ω) is a compact Kähler manifold of dimension n . For $(\varphi', \varphi'') \in \mathcal{P}_\omega \times \mathcal{P}_\omega$, Aubin-Yau functionals are defined by

$$(4.1) \quad \mathcal{I}_\omega^{\text{AY}}(\varphi', \varphi'') = \frac{1}{V_\omega} \int_X (\varphi'' - \varphi')(\omega_{\varphi'}^n - \omega_{\varphi''}^n),$$

$$(4.2) \quad \mathcal{J}_\omega^{\text{AY}}(\varphi', \varphi'') = -\mathcal{L}_\omega^M(\varphi', \varphi'') + \frac{1}{V_\omega} \int_X (\varphi'' - \varphi')\omega_{\varphi'}^n.$$

By definition, we have

$$(4.3) \quad \mathcal{J}_\omega^{\text{AY}}(\varphi', \varphi'') + \mathcal{J}_\omega^{\text{AY}}(\varphi'', \varphi') = \mathcal{I}_\omega^{\text{AY}}(\varphi', \varphi'') = \mathcal{I}_\omega^{\text{AY}}(\varphi'', \varphi').$$

For any $\varphi \in \mathcal{P}_\omega$ we set

$$(4.4) \quad \mathcal{I}_\omega^{\text{AY}}(\varphi) = \frac{1}{V_\omega} \int_X \varphi(\omega^n - \omega_\varphi^n),$$

$$(4.5) \quad \mathcal{J}_\omega^{\text{AY}}(\varphi) = \int_0^1 \frac{\mathcal{I}_\omega(s \cdot \varphi)}{s} ds = \frac{1}{V_\omega} \int_0^1 \int_X \varphi(\omega^n - \omega_{s \cdot \varphi}^n) ds.$$

It's clear that

$$(4.6) \quad \mathcal{I}_\omega^{\text{AY}}(\varphi) = \mathcal{I}_\omega^{\text{AY}}(0, \varphi), \quad \mathcal{J}_\omega^{\text{AY}}(\varphi) = \mathcal{J}_\omega^{\text{AY}}(0, \varphi).$$

By definition we have

$$\begin{aligned}
& \mathcal{J}_\omega^{\text{AY}}(\varphi) \\
&= \frac{1}{V_\omega} \int_0^1 ds \int_X \varphi(-\sqrt{-1} \partial \bar{\partial}(s \cdot \varphi)) \wedge \sum_{i=1}^{n-1} \omega^{n-1-i} \wedge \omega_{s \cdot \varphi}^i \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_0^1 s \cdot ds \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{i=0}^{n-1} \omega^{n-1-i} \wedge [\omega + s(\omega_\varphi - \omega)]^i \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_0^1 s \cdot ds \int_X \varphi \cdot \partial \bar{\partial} \varphi \sum_{i=0}^{n-1} \omega^{n-1-i} \wedge \sum_{j=0}^i \binom{i}{j} (1-s)^{i-j} s^j \omega^{i-j} \wedge \omega_\varphi^j \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{j=0}^{n-1} \sum_{i=j}^{n-1} \binom{i}{j} \omega^{n-1-j} \wedge \omega_\varphi^j \int_0^1 (1-s)^{i-j} s^{1+j} \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{j=0}^{n-1} \omega^{n-1-j} \wedge \omega_\varphi^j \cdot \sum_{i=j}^{n-1} \binom{i}{j} \cdot \frac{(i-j)! \cdot (j+1)!}{(i+2)!} \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{j=0}^{n-1} \omega^{n-1-j} \wedge \omega_\varphi^j \cdot \sum_{i=j}^{n-1} \frac{j+1}{(i+2)(i+1)} \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{j=0}^{n-1} \frac{n-j}{n+1} \omega^{n-1-j} \wedge \omega_\varphi^j
\end{aligned}$$

since

$$\sum_{i=j}^{n-1} \frac{1}{(i+2)(i+1)} = \sum_{i=j}^{n-1} \left(\frac{1}{i+1} - \frac{1}{i+2} \right) = \frac{1}{j+1} - \frac{1}{n+1}.$$

On the other hand,

$$\frac{n}{n+1} \mathcal{I}_\omega^{\text{AY}}(\varphi) = \frac{-\sqrt{-1}}{V_\omega} \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{i=0}^{n-1} \frac{n}{n+1} \omega^{n-1-i} \wedge \omega_\varphi^i$$

Hence

$$\begin{aligned}
(4.7) \quad & \frac{n}{n+1} \mathcal{I}_\omega^{\text{AY}}(\varphi) - \mathcal{J}_\omega^{\text{AY}}(\varphi) \\
&= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \sum_{j=1}^{n-1} \frac{j}{n+1} \omega^{n-1-j} \wedge \omega_\varphi^j.
\end{aligned}$$

Moreover,

$$\begin{aligned}
(4.8) \quad & (n+1) \mathcal{J}_\omega^{\text{AY}}(\varphi) - \mathcal{I}_\omega^{\text{AY}}(\varphi) \\
&= \frac{1}{V_\omega} \int_X \varphi \cdot -\sqrt{-1} \partial \bar{\partial} \varphi \wedge \sum_{j=0}^{n-1} (n-1-j) \omega^{n-1-j} \wedge \omega_\varphi^j.
\end{aligned}$$

Remark 4.1. Notice that formulae (4.7) and (4.8) are also valid when ω is non-Kähler.

4.2. Aubin-Yau functionals over compact complex surfaces. Let (X, g) be a compact complex manifold of the complex dimension n and ω be its associated real $(1, 1)$ -form. From Remark 4.1, we can formally use the notion $\mathcal{I}_\omega^{\text{AY}}$, $\mathcal{J}_\omega^{\text{AY}}$, and $\mathcal{L}_\omega^{\text{M}}$, but now ω may not be Kähler. Precisely, for any $\varphi \in \mathcal{P}_\omega$ we set

$$(4.9) \quad \mathcal{I}_{\omega|\bullet}^{\text{AY}}(\varphi) = \frac{1}{V_\omega} \int_X \varphi(\omega^n - \omega_\varphi^n),$$

$$(4.10) \quad \mathcal{J}_{\omega|\bullet}^{\text{AY}}(\varphi) = \int_0^1 \frac{\mathcal{I}_{\omega|\bullet}^{\text{AY}}(s \cdot \varphi)}{s} ds = \frac{1}{V_\omega} \int_0^1 \int_X \varphi(\omega^n - \omega_{s \cdot \varphi}^n) ds.$$

Hence

$$(4.11) \quad \begin{aligned} & \frac{n}{n+1} \mathcal{I}_{\omega|\bullet}^{\text{AY}}(\varphi) - \mathcal{J}_{\omega|\bullet}^{\text{AY}}(\varphi) \\ &= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \sum_{j=1}^{n-1} \frac{j}{n+1} \omega^{n-1-j} \wedge \omega_\varphi^j. \end{aligned}$$

Moreover,

$$(4.12) \quad \begin{aligned} & (n+1) \mathcal{J}_{\omega|\bullet}^{\text{AY}}(\varphi) - \mathcal{I}_{\omega|\bullet}^{\text{AY}}(\varphi) \\ &= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \sum_{j=0}^{n-1} (n-1-j) \omega^{n-1-j} \wedge \omega_\varphi^j. \end{aligned}$$

Restricting to compact complex surfaces and introducing two extra functionals on $\mathcal{P}_\omega$

$$(4.13) \quad \mathcal{A}_\omega(\varphi) := \frac{1}{2V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \omega \wedge \bar{\partial} \varphi),$$

$$(4.14) \quad \mathcal{B}_\omega(\varphi) := \frac{1}{2V_\omega} \int_X \varphi \cdot \sqrt{-1} \partial \omega \wedge \bar{\partial} \varphi,$$

(clearly $\overline{\mathcal{A}_\omega(\varphi)} = \mathcal{B}_\omega(\varphi)$), we define Aubin-Yau functionals as follows (Here constants a, b, c, d are determined later, and actually $a = b = c = d = 2$)

$$(4.15) \quad \mathcal{I}_\omega^{\text{AY}}(\varphi) := \mathcal{I}_{\omega|\bullet}^{\text{AY}}(\varphi) + a \mathcal{A}_\omega(\varphi) + b \mathcal{B}_\omega(\varphi),$$

$$(4.16) \quad \mathcal{J}_\omega^{\text{AY}}(\varphi) := -\mathcal{L}_\omega^{\text{M}}(\varphi) + \frac{1}{V_\omega} \int_X \varphi \cdot \omega^2 + c \mathcal{A}_\omega(\varphi) + d \mathcal{B}_\omega(\varphi).$$

Since

$$\begin{aligned} \mathcal{J}_{\omega|\bullet}^{\text{AY}}(\varphi) &= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \sum_{j=0}^1 \frac{2-j}{3} (\omega^{1-j} \wedge \omega_\varphi^j) \\ &= \frac{1}{V_\omega} \int_X \varphi (\omega - \omega_\varphi) \wedge \left(\frac{2}{3} \omega + \frac{1}{3} \omega_\varphi \right) \\ &= \frac{1}{V_\omega} \int_X \varphi \left(\frac{2}{3} \omega^2 - \frac{1}{3} \omega \wedge \omega_\varphi - \frac{1}{3} \omega_\varphi^2 \right), \end{aligned}$$

it follows that

$$(4.17) \quad \mathcal{J}_\omega^{\text{AY}}(\varphi) = \mathcal{J}_{\omega|\bullet}^{\text{AY}}(\varphi) + (c-1) \mathcal{A}_\omega(\varphi) + (d-1) \mathcal{B}_\omega(\varphi).$$

Using Remark 4.1 or the previous subsection, we deduce

$$\begin{aligned}
& \frac{2}{3}(\mathcal{I}_\omega^{\text{AY}}(\varphi) - a\mathcal{A}_\omega(\varphi) - b\mathcal{B}_\omega(\varphi)) - (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1)\mathcal{A}_\omega(\varphi) - (d-1)\mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{V_\omega} \int_X \varphi(-\sqrt{-1}\partial\bar{\partial}\varphi) \sum_{j=1}^1 \frac{j}{3} \omega^{1-j} \wedge \omega_\varphi^j \\
&= \frac{1}{V_\omega} \int_X \varphi \cdot \frac{1}{3} \omega_\varphi \wedge (-\sqrt{-1}\partial\bar{\partial}\varphi) = \frac{1}{V_\omega} \int_X \varphi \cdot \frac{1}{3} \omega_\varphi \wedge \sqrt{-1}\partial\bar{\partial}\varphi.
\end{aligned}$$

Thus the left hand side has two slightly different expressions. If we adopt the first one, we have

$$\begin{aligned}
& \frac{2}{3}(\mathcal{I}_\omega^{\text{AY}}(\varphi) - a\mathcal{A}_\omega(\varphi) - b\mathcal{B}_\omega(\varphi)) - (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1)\mathcal{A}_\omega(\varphi) - (d-1)\mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{3V_\omega} \int_X \sqrt{-1}\partial(\varphi \cdot \omega_\varphi) \wedge \bar{\partial}\varphi = \frac{1}{3V_\omega} \int_X \sqrt{-1}(\partial\varphi \wedge \omega_\varphi + \varphi \cdot \partial\omega) \wedge \bar{\partial}\varphi \\
&= \frac{1}{3V_\omega} \int_X \sqrt{-1}\partial\varphi \wedge \bar{\partial}\varphi \wedge \omega_\varphi - \frac{2}{3}\mathcal{A}_\omega(\varphi).
\end{aligned}$$

On the other hand, using the second expression gives

$$\begin{aligned}
& \frac{2}{3}(\mathcal{I}_\omega^{\text{AY}}(\varphi) - a\mathcal{A}_\omega(\varphi) - b\mathcal{B}_\omega(\varphi)) - (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1)\mathcal{A}_\omega(\varphi) - (d-1)\mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{3V_\omega} \int_X -\sqrt{-1}\bar{\partial}(\varphi \cdot \omega_\varphi) \wedge \partial\varphi = \frac{1}{3V_\omega} \int_X -\sqrt{-1}(\bar{\partial}\varphi \wedge \omega_\varphi + \varphi \cdot \bar{\partial}\omega) \wedge \partial\varphi \\
&= \frac{1}{3V_\omega} \int_X \sqrt{-1}\partial\varphi \wedge \bar{\partial}\varphi \wedge \omega_\varphi - \frac{2}{3}\mathcal{B}_\omega(\varphi).
\end{aligned}$$

Therefore

$$\begin{aligned}
& \frac{2}{3}(\mathcal{I}_\omega^{\text{AY}}(\varphi) - a\mathcal{A}_\omega(\varphi) - b\mathcal{B}_\omega(\varphi)) - (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1)\mathcal{A}_\omega(\varphi) - (d-1)\mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{3V_\omega} \int_X \sqrt{-1}\partial\varphi \wedge \bar{\partial}\varphi \wedge \omega_\varphi - \frac{\mathcal{A}_\omega(\varphi) + \mathcal{B}_\omega(\varphi)}{3},
\end{aligned}$$

or, equivalently,

$$(4.18) \quad \frac{2}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi) - \mathcal{J}_\omega^{\text{AY}}(\varphi) = \frac{1}{3V_\omega} \int_X \sqrt{-1}\partial\varphi \wedge \bar{\partial}\varphi \wedge \omega_\varphi$$

where we require

$$(4.19) \quad \frac{2}{3}a - (c-1) - \frac{1}{3} = 0 = \frac{2}{3}b - (d-1) - \frac{1}{3}.$$

Theorem 4.2. *For any $\varphi \in \mathcal{P}_\omega$, one has*

$$(4.20) \quad \frac{2}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi) - \mathcal{J}_\omega^{\text{AY}}(\varphi) \geq 0.$$

Using (4.12) yields

$$\begin{aligned}
& 3 \left(\mathcal{J}_\omega^{\text{AY}}(\varphi) - \frac{1}{2} \mathcal{A}_\omega(\varphi) - \frac{1}{2} \mathcal{B}_\omega(\varphi) \right) - (\mathcal{I}_\omega^{\text{AY}}(\varphi) - \mathcal{A}_\omega(\varphi) - \mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \sum_{j=0}^1 (1-j) \omega^{1-j} \wedge \omega_\varphi^j \\
&= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \omega \\
&= \frac{1}{V_\omega} \int_X (\varphi \cdot \omega) \wedge (-\sqrt{-1} \partial \bar{\partial} \varphi) = \frac{1}{V_\omega} \int_X (\varphi \cdot \omega) \wedge \sqrt{-1} \partial \bar{\partial} \varphi.
\end{aligned}$$

As the proof of Theorem 4.2, we have

$$\begin{aligned}
& 3 (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1) \mathcal{A}_\omega(\varphi) - (d-1) \mathcal{B}_\omega(\varphi)) - (\mathcal{I}_\omega^{\text{AY}}(\varphi) - a \mathcal{A}_\omega(\varphi) - b \mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{V_\omega} \int_X \sqrt{-1} \partial (\varphi \cdot \omega) \wedge \bar{\partial} \varphi = \frac{1}{V_\omega} \int_X \sqrt{-1} (\partial \varphi \wedge \omega + \varphi \cdot \partial \omega) \wedge \bar{\partial} \varphi \\
&= \frac{1}{V_\omega} \int_X \sqrt{-1} \partial \varphi \wedge \bar{\partial} \varphi \wedge \omega - 2 \mathcal{A}_\omega(\varphi)
\end{aligned}$$

and

$$\begin{aligned}
& 3 (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1) \mathcal{A}_\omega(\varphi) - (d-1) \mathcal{B}_\omega(\varphi)) - (\mathcal{I}_\omega^{\text{AY}}(\varphi) - a \mathcal{A}_\omega(\varphi) - b \mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{V_\omega} \int_X -\sqrt{-1} \bar{\partial} (\varphi \cdot \omega) \wedge \partial \varphi = \frac{1}{V_\omega} \int_X -\sqrt{-1} (\bar{\partial} \varphi \wedge \omega + \varphi \cdot \bar{\partial} \omega) \wedge \partial \varphi \\
&= \frac{1}{V_\omega} \int_X \sqrt{-1} \bar{\partial} \varphi \wedge \partial \varphi \wedge \omega - 2 \mathcal{B}_\omega(\varphi).
\end{aligned}$$

Hence

$$\begin{aligned}
& 3 (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1) \mathcal{A}_\omega(\varphi) - (d-1) \mathcal{B}_\omega(\varphi)) - (\mathcal{I}_\omega^{\text{AY}}(\varphi) - a \mathcal{A}_\omega(\varphi) - b \mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{V_\omega} \int_X \sqrt{-1} \partial \varphi \wedge \bar{\partial} \varphi \wedge \omega - (\mathcal{A}_\omega(\varphi) + \mathcal{B}_\omega(\varphi)).
\end{aligned}$$

Equivalently,

$$(4.21) \quad 3 \mathcal{J}_\omega^{\text{AY}}(\varphi) - \mathcal{I}_\omega^{\text{AY}}(\varphi) = \frac{1}{V_\omega} \int_X \sqrt{-1} \partial \varphi \wedge \bar{\partial} \varphi \wedge \omega,$$

where we also require

$$(4.22) \quad 3(c-1) - a - 1 = 0 = 3(d-1) - b - 1.$$

Theorem 4.3. *For any $\varphi \in \mathcal{P}_\omega$, one has*

$$(4.23) \quad 3 \mathcal{J}_\omega^{\text{AY}}(\varphi) - \mathcal{I}_\omega^{\text{AY}}(\varphi) \geq 0.$$

Combining (4.19) and (4.22) we obtain the value of those constants:

$$a = b = c = d = 2.$$

Corollary 4.4. *For any compact complex surface (X, g) and any real-valued smooth function $\varphi \in \mathcal{P}_\omega$, we have*

$$\begin{aligned} \frac{1}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi) &\leq \mathcal{J}_\omega^{\text{AY}}(\varphi) \leq \frac{2}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi), \\ \frac{3}{2}\mathcal{J}_\omega^{\text{AY}}(\varphi) &\leq \mathcal{I}_\omega(\varphi) \leq 3\mathcal{J}_\omega^{\text{AY}}(\varphi), \\ \frac{1}{2}\mathcal{J}_\omega^{\text{AY}}(\varphi) &\leq \frac{1}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi) \leq \mathcal{I}_\omega^{\text{AY}}(\varphi) - \mathcal{J}_\omega^{\text{AY}}(\varphi) \leq \frac{2}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi) \leq 2\mathcal{J}_\omega^{\text{AY}}(\varphi), \end{aligned}$$

where ω is its associated real $(1, 1)$ -form.

REFERENCES

1. Chen, X. X., Tian, G., *Ricci flow on Kähler-Einstein surfaces*, Invent. Math. **147**(2002), no.3, 487–544.
2. Dinew, S., Kolodziej, S., *Pluripotential estimates on compact Hermitian manifolds*, preprint, arXiv: 0910.3937, 2009.
3. Gauduchon, P., *Sur la 1-forme de torsion d'une variété hermitienne compacte*, Math. Ann. **267** (1984), 495–518.
4. Guan, B., Li, Q., *Complex Monge-Ampère equations on Hermitian manifold*, preprint, arXiv: 0906.3548, 2009.
5. Lübke, M., Teleman, A., *The Kobayashi-Hitchin correspondence*, World Scientific, 1995.
6. Mabuchi, T., *K-energy maps integrating Futaki invariants*, Tohoku Math. Journ., **38**(1986), 575–593.
7. Phong, D.H., Sturm, J., *Lectures on stability and constant scalar curvature*, preprint, arXiv: 0801.4179, 2008.
8. Siu, Y.T., *Every K3-surface is Kähler*, Invent. Math., **73** (1983), 139–150.
9. Tosatti, V., Weinkove, B., *Estimates for the complex Monge-Ampère equation on Hermitian and balanced manifolds*, preprint, arXiv: 0909.4496, 2009.
10. Tosatti, V., Weinkove, B., *The complex Monge-ampère equation on compact Hermitian manifolds*, preprint, arXiv: 0910.1390, 2009.
11. Yau, S.-T., *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I*, Comm. Pure. Appl. Math., **31** (1978), no.3, 339–411.
12. Yau, S.-T., *Review of geometry and analysis*, Kodaira's issue. Asian J. Math. **4**(2000), no.1, 235–278.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA 02138
E-mail address: yili@math.harvard.edu

MABUCHI AND AUBIN-YAU FUNCTIONALS OVER COMPLEX SURFACES

YI LI

*This paper is dedicated to my colleague and friend Lin Chen
who has passed away in an accident.*

ABSTRACT. In this note we construct Mabuchi $\mathcal{L}_\omega^M$ functional and Aubin-Yau functionals $\mathcal{I}_\omega^{AY}, \mathcal{J}_\omega^{AY}$ on any compact complex surfaces, and establish a number of properties. Our construction coincides with the original one in the Kähler case.

CONTENTS

1. Introduction	1
2. Err_ω map on complex manifolds	3
2.1. Err_ω map on compact complex manifolds	3
2.2. Vanishing property of Err_ω map on compact complex surface	6
3. Mabuchi $\mathcal{L}_\omega^M$ functional on complex surfaces	7
3.1. Mabuchi $\mathcal{L}_\omega^M$ functional on compact Kähler manifolds	7
3.2. Mabuchi $\mathcal{L}_\omega^M$ functional on compact complex surfaces	7
4. Aubin-Yau functionals on compact complex surfaces	11
4.1. Aubin-Yau functionals on compact Kähler manifolds	11
4.2. Aubin-Yau functionals over compact complex surfaces	13
References	16

1. INTRODUCTION

Let (X, ω) be a compact Kähler manifold of the complex dimension n . It's known that the volume V_ω depends only on the Kähler class of ω , namely,

$$\int_X (\omega + \sqrt{-1}\partial\bar{\partial}\varphi)^n = \int_X \omega^n$$

for any real-valued smooth function φ with $\omega_\varphi := \omega + \sqrt{-1}\partial\bar{\partial}\varphi > 0$, because of closedness of ω .

If (X, g) is a compact Hermitian manifold of the complex dimension n , the same result does not hold in general. In the Section 2, we consider a function to describe such a phenomena, i.e., we define

$$\text{Err}_\omega(\varphi) := \int_X \omega^n - \int_X \omega_\varphi^n,$$

where ω is its associated real $(1, 1)$ -form. The main result ¹ is

Proposition 1.1. *Let (X, g) be a compact Hermitian manifold of the complex dimension n and ω its associated real $(1, 1)$ -form. If $\partial\bar{\partial}(\omega^k) = 0$ for $k = 1, 2$, then*

$$\int_X \omega_\varphi^n = \int_X \omega^n =: V_\omega$$

for any real-valued function $\varphi \in C^\infty(X)_\mathbb{R}$ with $\omega_\varphi := \omega + \sqrt{-1}\partial\bar{\partial}\varphi > 0$.

In Kähler geometry, energy functionals, such as Mabuchi K -energy functional [6], Aubin-Yau energy functionals [7], and Chen-Tian energy functionals [1], play an important role in studying Kähler-Einstein metrics and constant scalar curvatures. When I was in Yau's Seminar, I asked myself that can we define energy functionals on compact complex manifolds? This is one motivation to write down this note. Another motivation comes from a question in S.-T. Yau's survey [12], that find necessary and sufficient conditions for a complex manifold to admit a Kähler structure. When $n = 2$, it was settled by Siu [8] or see [5]: A compact complex surface is Kähler if and only if its first Betti number is even. In the second part of this note we construct Mabuchi $\mathcal{L}_\omega^M$ functional and Aubin-Yau functionals $\mathcal{I}_\omega^{AY}, \mathcal{J}_\omega^{AY}$ on any compact complex surface.

Let (X, g) be a compact Hermitian manifold of the complex dimension 2 and ω be its associated real $(1, 1)$ -form. Let

$$\mathcal{P}_\omega := \{\varphi \in C^\infty(X)_\mathbb{R} | \omega_\varphi := \omega + \sqrt{-1}\partial\bar{\partial}\varphi > 0\}.$$

For any $\varphi', \varphi'' \in \mathcal{P}_\omega$, we define

$$\begin{aligned} \mathcal{L}_\omega^M(\varphi', \varphi'') &:= \frac{1}{V_\omega} \int_0^1 \int_X \dot{\varphi}_t \cdot \omega_{\varphi_t}^2 dt \\ &\quad - \frac{1}{V_\omega} \int_0^1 \int_X \sqrt{-1}\partial\omega \wedge (\bar{\partial}\dot{\varphi}_t \cdot \varphi_t) dt \\ &\quad + \frac{1}{V_\omega} \int_0^1 \int_X \sqrt{-1}\bar{\partial}\omega \wedge (\partial\dot{\varphi}_t \cdot \varphi_t) dt \end{aligned}$$

where φ_t is any smooth path in $\mathcal{P}_\omega$ from φ' to φ'' . Also we set

$$\mathcal{L}_\omega^M(\varphi) := \mathcal{L}_\omega^M(0, \varphi).$$

Theorem 1.2. *The functional $\mathcal{L}_\omega^M(\varphi', \varphi'')$ is independent of the choice of the smooth path $\{\varphi_t\}_{0 \leq t \leq 1}$ and satisfies the 1-cocycle condition. In particular*

$$\begin{aligned} \mathcal{L}_\omega^M(\varphi) &= \frac{1}{3V_\omega} \int_X \varphi(\omega^2 + \omega \wedge \omega_\varphi + \omega_\varphi^2) \\ &\quad + \frac{1}{2V_\omega} \int_X \varphi[-\sqrt{-1}\partial\omega \wedge \bar{\partial}\varphi + \sqrt{-1}\bar{\partial}\omega \wedge \partial\varphi]. \end{aligned}$$

Moreover, for any $\varphi \in \mathcal{P}_\omega$ and any constant $C \in \mathbb{R}$, we have

$$\mathcal{L}_\omega^M(\varphi, \varphi + C) = C \cdot \left(1 - \frac{\text{Err}_\omega(\varphi)}{V_\omega}\right);$$

for any $\varphi_1, \varphi_2 \in \mathcal{P}_\omega$ and any constant $C \in \mathbb{R}$, we have

$$\mathcal{L}_\omega^M(\varphi_1, \varphi_2 + C) = \mathcal{L}_\omega^M(\varphi_1, \varphi_2) + C \cdot \left(1 - \frac{\text{Err}_\omega(\varphi)}{V_\omega}\right).$$

¹V. Tosatti told the author that the same result has implicitly contained in [10] and [2].

In Section 4, we construct Aubin-Yau functionals on compact complex surfaces. Let (X, g) be a compact Hermitian manifold of the complex dimension 2 and ω be its associated real $(1, 1)$ -form. Set

$$\begin{aligned}\mathcal{A}_\omega(\varphi) &:= \frac{1}{2V_\omega} \int_X \varphi \cdot -\sqrt{-1} \partial \omega \wedge \bar{\partial} \varphi, \\ \mathcal{B}_\omega(\varphi) &:= \frac{1}{2V_\omega} \int_X \varphi \cdot \sqrt{-1} \bar{\partial} \omega \wedge \partial \varphi.\end{aligned}$$

We define

$$\begin{aligned}\mathcal{I}_\omega^{\text{AY}}(\varphi) &:= \frac{1}{V_\omega} \int_X \varphi (\omega^2 - \omega_\varphi^2) + 2\mathcal{A}_\omega(\varphi) + 2\mathcal{B}_\omega(\varphi), \\ \mathcal{J}_\omega^{\text{AY}}(\varphi) &:= \frac{1}{V_\omega} \int_0^1 \int_X \varphi (\omega^2 - \omega_{s \cdot \varphi}^2) ds + \mathcal{A}_\omega(\varphi) + \mathcal{B}_\omega(\varphi).\end{aligned}$$

Theorem 1.3. *For any compact Hermitian manifold (X, g) of the complex dimension 2, we have*

$$\frac{1}{3} \mathcal{I}_\omega^{\text{AY}}(\varphi) \leq \mathcal{J}_\omega^{\text{AY}}(\varphi) \leq \frac{2}{3} \mathcal{I}_\omega^{\text{AY}}(\varphi)$$

for any $\varphi \in \mathcal{P}_\omega$, where ω is its associated real $(1, 1)$ -form.

We hope this exposition will give some ideas to study Yau's problem. The author are concerning the construction of those functionals on higher dimensional compact complex manifolds.

Acknowledgements. The author would like to thank Valentino Tosatti who read this note and pointed out a serious mistake in the first version. Furthermore, I also thank Chen-Yu Chi and Ming-Tao Chuan for useful discussion.

2. Err $_\omega$ MAP ON COMPLEX MANIFOLDS

Let (X, g) be a compact Hermitian manifold of the complex dimension n and write the associated real $(1, 1)$ -form as

$$\omega = \sqrt{-1} \cdot g_{i\bar{j}} \cdot dz^i \wedge d\bar{z}^j.$$

Let $\mathcal{P}_\omega$ be the space of all real-valued smooth functions $\varphi \in C^\infty(X)_\mathbb{R}$ so that $\omega + \sqrt{-1} \partial \bar{\partial} \varphi$ is positive definite on X :

$$\mathcal{P}_\omega := \{\varphi \in C^\infty(X)_\mathbb{R} \mid \omega + \sqrt{-1} \partial \bar{\partial} \varphi > 0\}.$$

Also we set

$$\mathcal{P}_\omega^0 := \left\{ \varphi \in \mathcal{P}_\omega \mid \sup_X \varphi = 0 \right\}.$$

2.1. Err $_\omega$ map on compact complex manifolds. To such a function $\varphi \in \mathcal{P}_\omega$ we associate the quantity

$$(2.1) \quad V_\omega(\varphi) := \int_X \omega_\varphi^n$$

the volume of X with respect to φ . In particular we set

$$(2.2) \quad V_\omega := V_\omega(0) = \int_X \omega^n.$$

For X being Kähler, we have $V_\omega = V_\omega(\varphi)$ for any $\varphi \in \mathcal{P}_\omega$. In the non-Kähler case, it's not in general true. Hence it is reasonable to define

$$(2.3) \quad \text{Err}_\omega(\varphi) := V_\omega - V_\omega(\varphi) = \int_X \omega^n - \int_X \omega_\varphi^n.$$

A natural question is when does $\text{Err}_\omega(\varphi)$ vanish for any/a $\varphi \in \mathcal{P}_\omega$? Clearly there exists a smooth real-valued function $\varphi_0 \equiv 0 \in \mathcal{P}_\omega$ such that

$$\omega_{\varphi_0}^n = \omega^n, \quad \sup_X \varphi_0 = 0,$$

hence

$$(2.4) \quad \text{Err}_\omega(\varphi_0) = \int_X \omega^n - \int_X \omega^n = 0.$$

This gives us some information about $\text{Err}_\omega(\varphi)$ and motivates us to consider

$$(2.5) \quad \text{SupErr}_\omega := \sup_{\varphi \in \mathcal{P}_\omega^0} (\text{Err}_\omega(\varphi)), \quad \text{InfErr}_\omega := \inf_{\varphi \in \mathcal{P}_\omega^0} (\text{Err}_\omega(\varphi)).$$

In any case, one has

$$(2.6) \quad \text{InfErr}_\omega \leq 0 \leq \text{SupErr}_\omega \leq \int_X \omega^n.$$

It's interesting to find some conditions to guarantee that the equalities hold. To study this behavior of Err_ω we consider the following several natural conditions on ω :

• **Condition 1.1:**

$$(2.7) \quad \sqrt{-1}\partial\omega \wedge \bar{\partial}\omega \quad \text{and} \quad \sqrt{-1}\partial\bar{\partial}\omega \quad \text{are non-negative}$$

• **Condition 1.2:**

$$(2.8) \quad \sqrt{-1}\partial\omega \wedge \bar{\partial}\omega \quad \text{and} \quad \sqrt{-1}\partial\bar{\partial}\omega \quad \text{are non-positive}$$

• **Condition 2:**

$$(2.9) \quad \partial\bar{\partial}(\omega^k) = 0, \quad k = 1, 2.$$

• **Condition 3:**

$$(2.10) \quad d(\omega^{n-1}) = 0.$$

• **Condition 4:**

$$(2.11) \quad \partial\bar{\partial}(\omega^{n-1}) = 0.$$

Remark 2.1. Condition 2 was appeared in [4] as a sufficient condition to solving the complex Monge-Ampère equation on Hermitian manifolds. The metric satisfying the third condition is called a balanced metric, which naturally appears in string theory (V. Tosatti and B. Wenkove [9] solved the complex Monge-Ampère equation on Hermitian manifolds with balanced metrics; later, they [10] dropped off the balanced condition.); When $n = 2$, this condition is indeed the Kähler condition. A metric satisfying Condition 4 is referred to be a Gauduchon metric, and a theorem of Gauduchon [3] shows that there exists a Gauduchon metric on every compact Hermitian manifold. Notice that Condition 3 implies condition 4, and Condition 2 is equivalent to $\partial\bar{\partial}\omega = 0 = \partial\omega \wedge \bar{\partial}\omega$. In particular, Condition 2 implies Condition 1.1 and Condition 1.2. In our case $n = 2$, Condition 2 is equivalent to Condition 4.

Remark 2.2. For any two forms α of degree $|\alpha|$ and β of degree $|\beta|$, we have

$$\alpha \wedge \beta = (-1)^{|\alpha| \cdot |\beta|} \beta \wedge \alpha.$$

Also we have

$$d(\alpha \wedge \beta) = d\alpha \wedge \beta + (-1)^{|\alpha|} \alpha \wedge d\beta;$$

according to types, one deduces

$$\begin{aligned} \partial(\alpha \wedge \beta) &= \partial\alpha \wedge \beta + (-1)^{|\alpha|} \alpha \wedge \partial\beta, \\ \bar{\partial}(\alpha \wedge \beta) &= \bar{\partial}\alpha \wedge \beta + (-1)^{|\alpha|} \alpha \wedge \bar{\partial}\beta. \end{aligned}$$

Moreover, if $|\alpha| + |\beta| = 2n - 1$, then

$$\begin{aligned} \int_X \alpha \wedge \partial\beta &= (-1)^{|\beta|} \int_X \partial\alpha \wedge \beta = -(-1)^{|\alpha|} \int_X \partial\alpha \wedge \beta, \\ \int_X \alpha \wedge \bar{\partial}\beta &= (-1)^{|\beta|} \int_X \bar{\partial}\alpha \wedge \beta = -(-1)^{|\alpha|} \int_X \bar{\partial}\alpha \wedge \beta. \end{aligned}$$

Those formulae are useful in our computations.

Theorem 2.3. (i) If ω satisfies Condition 1.1, then

$$(2.12) \quad \text{InfErr}_\omega = 0.$$

(ii) Correspondingly, if ω satisfies Condition 1.2, then

$$(2.13) \quad \text{SupErr}_\omega = 0.$$

(iii) In particular $\text{SupErr}_\omega = \text{InfErr}_\omega = 0$ provided that ω satisfies Condition 2.

Proof. (i) we knew that $\text{Err}_\omega(\varphi_0) = 0$ for some $\varphi_0 \in \mathcal{P}_\omega^0$. To achieve the argument, we need only to show that $\text{Err}_\omega(\varphi) \geq 0$ for each function $\varphi \in \mathcal{P}_\omega^0$. By definition we have

$$\begin{aligned} &\text{Err}_\omega(\varphi) \\ &= - \int_X \omega_\varphi^n + \int_X \omega^n = \int_X -\sqrt{-1} \partial \bar{\partial} \varphi \wedge \sum_{i=0}^{n-1} \omega_\varphi^i \wedge \omega^{n-1-i} \\ &= \sum_{i=0}^{n-1} \int_X \omega_\varphi^i \wedge \omega^{n-1-i} \wedge (-\sqrt{-1} \partial \bar{\partial} \varphi) = \sum_{i=0}^{n-1} \int_X \sqrt{-1} \partial (\omega_\varphi^i \wedge \omega^{n-1-i}) \wedge \bar{\partial} \varphi \\ &= \sum_{i=0}^{n-1} \int_X [i \cdot \omega_\varphi^{i-1} \wedge \partial \omega \wedge \omega^{n-1-i} + \omega_\varphi^i \wedge (n-1-i) \omega^{n-2-i} \wedge \partial \omega] \wedge \sqrt{-1} \bar{\partial} \varphi \\ &= \sum_{i=0}^{n-1} \int_X [i \cdot \omega_\varphi^{i-1} \wedge \omega^{n-1-i} + (n-1-i) \omega_\varphi^i \wedge \omega^{n-2-i}] \wedge \partial \omega \wedge \sqrt{-1} \bar{\partial} \varphi \\ &= \sum_{i=0}^{n-1} \int_X \sqrt{-1} [\varphi \cdot (i \cdot \omega_\varphi^{i-1} \wedge \omega^{n-1-i} + (n-1-i) \cdot \omega_\varphi^i \wedge \omega^{n-2-i}) \wedge \bar{\partial} \partial \omega \\ &\quad + \varphi \cdot \bar{\partial} (i \cdot \omega_\varphi^{i-1} \wedge \omega^{n-1-i} + (n-1-i) \omega_\varphi^i \wedge \omega^{n-2-i}) \wedge \partial \omega] \\ &= \sum_{i=0}^{n-1} (I_i + II_i) \end{aligned}$$

where

$$\begin{aligned} I_i &= \int_X \varphi \cdot [i \cdot \omega_\varphi^{i-1} \wedge \omega^{n-1-i} + (n-1-i)\omega_\varphi^i \wedge \omega^{n-2-i}] \wedge (-\sqrt{-1}\partial\bar{\partial}\omega), \\ II_i &= \int_X \varphi \cdot \sqrt{-1}\partial\bar{\partial}[i \cdot \omega_\varphi^{i-1} \wedge \omega^{n-1-i} + (n-1-i)\omega_\varphi^i \wedge \omega^{n-2-i}] \wedge \partial\omega. \end{aligned}$$

Since $\sqrt{-1}\partial\bar{\partial}\omega \geq 0$ and $\varphi \leq 0$ on X , the first term I_i is non-negative. Applying the integration by parts to II_i , we deduce

$$\begin{aligned} II_i &= \int_X \varphi [i(i-1)\omega_\varphi^{i-2} \wedge \bar{\partial}\omega \wedge \omega^{n-1-i} + i(n-1-i)\omega_\varphi^{i-1} \wedge \omega^{n-2-i} \wedge \bar{\partial}\omega \\ &\quad + i(n-1-i)\omega_\varphi^{i-1} \wedge \bar{\partial}\omega \wedge \omega^{n-2-i} \\ &\quad + (n-1-i)(n-2-i)\omega_\varphi^i \wedge \omega^{n-3-i} \wedge \bar{\partial}\omega] \wedge \sqrt{-1}\partial\omega \\ &= \int_X \varphi \cdot \omega_\varphi^{i-2} \wedge \omega^{n-3-i} \wedge [i(i-1)\omega^2 + 2i(n-1-i)\omega_\varphi \wedge \omega \\ &\quad + (n-1-i)(n-2-i)\omega_\varphi^2] \wedge (-\sqrt{-1}\partial\omega \wedge \bar{\partial}\omega). \end{aligned}$$

Since $\sqrt{-1}\partial\omega \wedge \bar{\partial}\omega$ is non-negative and φ is non-positive, it follows that $II_i \geq 0$. Thus $\text{Err}_\omega(\varphi) \geq 0$ for each $\varphi \in \mathcal{P}_\omega^0$ and therefore $\text{InfErr}_\omega = 0$.

(ii) If ω satisfies Condition 1.2, the above proceeding gives that $\text{Err}_\omega(\varphi) \leq 0$ for each $\varphi \in \mathcal{P}_\omega^0$, i.e., $\text{SupErr}_\omega \leq 0$. Hence $\text{SupErr}_\omega = 0$.

(iii) It's an immediate consequence of (i) and (ii). $\square$

Corollary 2.4. *If ω satisfies Condition 2, then $\text{Err}_\omega(\varphi) = 0$ for any $\varphi \in \mathcal{P}_\omega^0$. Equivalently, in this case, the number $V_\omega(\varphi) = \int_X \omega_\varphi^n$ does not depend on the choice of $\varphi \in \mathcal{P}_\omega$ and equals $V_\omega = \int_X \omega^n$.*

2.2. Vanishing property of Err_ω map on compact complex surface. Let (X, g) be a Hermitian manifold of the complex dimension n and let ω_g be its associated real $(1, 1)$ -form. We say that g is a Gauduchon metric if $\partial\bar{\partial}(\omega_g^{n-1}) = 0$.

We recall a theorem of Gauduchon [3] or see Remark 2.1.

Theorem 2.5. (Gauduchon, 1984) *If X is a compact complex manifold of the complex dimension n , then in the conformal class of every Hermitian metric g there exists a Gauduchon metric g_G , i.e., there is a positive function $\varphi \in C^\infty(X)_\mathbb{R}$ such that $g_G := \varphi \cdot g$ is Gauduchon. If X is connected and $n \geq 2$, then g_G is unique up to a positive constants.*

Using the existence of Gauduchon metric, we can prove the following

Theorem 2.6. *Let (X, g) be a compact complex surface with Hermitian metric g and let ω_G be its associated Gauduchon metric. Then*

$$(2.14) \quad \text{Err}_{\omega_G}(\varphi) = 0$$

for all $\varphi \in \mathcal{P}_{\omega_G}$.

Proof. In what follows, we omit the subscript G and write ω_G as ω ; since ω is a Gauduchon metric, it follows that $\partial\bar{\partial}\omega = 0$. In the case of $n = 2$, we have

$$\begin{aligned} \text{Err}_\omega(\varphi) &= \int_X \omega^2 - \int_X \omega_\varphi^2 = \int_X (\omega - \omega_\varphi)(\omega + \omega_\varphi) \\ &= \int_X (\omega + \omega_\varphi) \wedge -\sqrt{-1}\partial\bar{\partial}\varphi \\ &= \int_X -\sqrt{-1}\partial\bar{\partial}(\omega + \omega_\varphi) \cdot \varphi \\ &= \int_X -2\sqrt{-1}\partial\bar{\partial}\omega \cdot \varphi = 0. \end{aligned}$$

Hence the theorem follows. $\square$

3. MABUCHI $\mathcal{L}_\omega^M$ FUNCTIONAL ON COMPLEX SURFACES

3.1. Mabuchi $\mathcal{L}_\omega^M$ functional on compact Kähler manifolds. Suppose that (X, ω) is a compact Kähler manifold of the complex dimension n . For any pair $(\varphi', \varphi'') \in \mathcal{P}_\omega \times \mathcal{P}_\omega$ we define

$$\mathcal{L}_\omega^M : \mathcal{P}_\omega \times \mathcal{P}_\omega \longrightarrow \mathbb{R}$$

as follows:

$$(3.1) \quad \mathcal{L}_\omega^M(\varphi', \varphi'') := \frac{1}{V_\omega} \int_0^1 \int_X \dot{\varphi}_t \cdot \omega_{\varphi_t}^n \cdot dt$$

where $\{\varphi_t : 0 \leq t \leq 1\}$ is any smooth path in $\mathcal{P}_\omega$ such that $\varphi_0 = \varphi'$ and $\varphi_1 = \varphi''$. For any $\varphi \in \mathcal{P}_\omega$ we set

$$(3.2) \quad \mathcal{L}_\omega^M(\varphi) := \mathcal{L}_\omega^M(0, \varphi).$$

Mabuchi [6] showed that the functional (3.1) is well-defined, and hence we can explicitly write down $\mathcal{L}_\omega^M(\varphi)$.

In this section we extend Mabuchi $\mathcal{L}_\omega^M$ functional to any compact complex surface by adding two extra terms on the right hand side of (3.1).

3.2. Mabuchi $\mathcal{L}_\omega^M$ functional on compact complex surfaces. Suppose now that (X, g) is a compact complex surface and ω be its associated real $(1, 1)$ -form. Let $\varphi', \varphi'' \in \mathcal{P}_\omega$ and $\{\varphi_t\}_{0 \leq t \leq 1}$ be a smooth path in $\mathcal{P}_\omega$ from φ' to φ'' .

Let

$$(3.3) \quad \mathcal{L}_\omega^0(\varphi', \varphi'') := \frac{1}{V_\omega} \int_0^1 \int_X \dot{\varphi}_t \cdot \omega_{\varphi_t}^2 \cdot dt.$$

Set

$$(3.4) \quad \psi(s, t) := s \cdot \varphi_t, \quad 0 \leq s \leq 1, \quad 0 \leq t \leq 1.$$

Consider a 1-form on $[0, 1] \times [0, 1]$

$$(3.5) \quad \Psi^0 := \left(\int_X \frac{\partial \psi}{\partial s} \cdot \omega_\psi^2 \right) ds + \left(\int_X \frac{\partial \psi}{\partial t} \cdot \omega_\psi^2 \right) dt.$$

Taking differential on Ψ^0 , we have

$$d\Psi^0 = I^0 \cdot dt \wedge ds$$

where

$$(3.6) \quad I^0 = \int_X \frac{\partial}{\partial t} \left(\frac{\partial \psi}{\partial s} \cdot \omega_\psi^2 \right) - \int_X \frac{\partial}{\partial s} \left(\frac{\partial \psi}{\partial t} \cdot \omega_\psi^2 \right).$$

Directly computing shows

$$\begin{aligned} I^0 &= \int_X \left[\frac{\partial^2 \psi}{\partial t \partial s} \cdot \omega_\psi^2 + \frac{\partial \psi}{\partial s} \cdot 2 \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \right] \\ &\quad - \int_X \left[\frac{\partial^2 \psi}{\partial s \partial t} \cdot \omega_\psi^2 + \frac{\partial \psi}{\partial t} \cdot 2 \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \right] \\ &= \int_X 2 \frac{\partial \psi}{\partial s} \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) - \int_X 2 \frac{\partial \psi}{\partial t} \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right). \end{aligned}$$

In the following we deduce two slightly different formulae of I^0 . The first one is

$$\begin{aligned} I^0 &= \int_X 2 \frac{\partial \psi}{\partial s} \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) + \int_X 2 \frac{\partial \psi}{\partial t} \cdot \omega_\psi \wedge \sqrt{-1} \partial \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \\ &= \int_X -2 \sqrt{-1} \partial \left(\frac{\partial \psi}{\partial s} \cdot \omega_\psi \right) \wedge \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \\ &\quad + \int_X -2 \sqrt{-1} \bar{\partial} \left(\frac{\partial \psi}{\partial t} \cdot \omega_\psi \right) \wedge \partial \left(\frac{\partial \psi}{\partial s} \right) \\ &= \int_X -2 \sqrt{-1} \left[\partial \left(\frac{\partial \psi}{\partial s} \right) \wedge \omega_\psi + \frac{\partial \psi}{\partial s} \cdot \partial \omega \right] \wedge \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \\ &\quad + \int_X -2 \sqrt{-1} \left[\bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \wedge \omega_\psi + \frac{\partial \psi}{\partial t} \cdot \bar{\partial} \omega \right] \wedge \partial \left(\frac{\partial \psi}{\partial s} \right) \\ &= \int_X -2 \sqrt{-1} \frac{\partial \psi}{\partial s} \cdot \partial \omega \wedge \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) + \int_X -2 \sqrt{-1} \frac{\partial \psi}{\partial t} \cdot \bar{\partial} \omega \wedge \partial \left(\frac{\partial \psi}{\partial s} \right) \\ &= \int_X 2 \sqrt{-1} \frac{\partial \psi}{\partial s} \cdot \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \wedge \partial \omega + \int_X 2 \sqrt{-1} \frac{\partial \psi}{\partial t} \cdot \partial \left(\frac{\partial \psi}{\partial s} \right) \wedge \bar{\partial} \omega. \end{aligned}$$

Similarly, we have

$$I^0 = \int_X -2 \sqrt{-1} \frac{\partial \psi}{\partial s} \cdot \partial \left(\frac{\partial \psi}{\partial t} \right) \wedge \bar{\partial} \omega + \int_X -2 \sqrt{-1} \frac{\partial \psi}{\partial t} \cdot \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \wedge \partial \omega.$$

Next, we define

$$(3.7) \quad \mathcal{L}_\omega^1(\varphi', \varphi'') = \frac{1}{V_\omega} \int_0^1 \int_X a_2 \cdot \partial \omega \wedge (\bar{\partial} \dot{\varphi}_t \cdot \varphi_t) dt,$$

$$(3.8) \quad \mathcal{L}_\omega^2(\varphi', \varphi'') = \frac{1}{V_\omega} \int_0^1 \int_X b_2 \cdot \bar{\partial} \omega \wedge (\partial \dot{\varphi}_t \cdot \varphi_t) dt.$$

Here we require $\bar{a}_2 = b_2$, and a_2, b_2 are determined later. As before, consider

$$\begin{aligned} \Psi^1 &= \left[\int_X a_2 \partial \omega \wedge \left(\bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \cdot \psi \right) \right] ds + \left[\int_X a_2 \partial \omega \wedge \left(\bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \cdot \psi \right) \right] dt, \\ \Psi^2 &= \left[\int_X b_2 \bar{\partial} \omega \wedge \left(\partial \left(\frac{\partial \psi}{\partial s} \right) \cdot \psi \right) \right] ds + \left[\int_X b_2 \bar{\partial} \omega \wedge \left(\partial \left(\frac{\partial \psi}{\partial t} \right) \cdot \psi \right) \right] dt. \end{aligned}$$

Therefore

$$d\Psi^1 = I^1 \cdot dt \wedge ds$$

where

$$(3.9) \quad \begin{aligned} I^1 &= \int_X a_2 \frac{\partial}{\partial t} \left[\partial \omega \wedge \left(\bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \cdot \psi \right) \right] \\ &\quad - \int_X a_2 \frac{\partial}{\partial s} \left[\partial \omega \wedge \left(\bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \cdot \psi \right) \right]. \end{aligned}$$

Dividing I^1 by a_2 yields

$$\begin{aligned} \frac{I^1}{a_2} &= \int_X -\frac{\partial}{\partial t} \left[\left(\bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \cdot \psi \right) \wedge \partial \omega \right] + \int_X \frac{\partial}{\partial s} \left[\left(\bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \cdot \psi \right) \wedge \partial \omega \right] \\ &= \int_X - \left[\bar{\partial} \left(\frac{\partial^2 \psi}{\partial t \partial s} \right) \cdot \psi + \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \cdot \frac{\partial \psi}{\partial t} \right] \wedge \partial \omega \\ &\quad + \int_X \left[\bar{\partial} \left(\frac{\partial^2 \psi}{\partial s \partial t} \right) \cdot \psi + \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \cdot \frac{\partial \psi}{\partial s} \right] \wedge \partial \omega \\ &= \int_X -\frac{\partial \psi}{\partial t} \cdot \bar{\partial} \left(\frac{\partial \psi}{\partial s} \right) \wedge \partial \omega + \int_X \frac{\partial \psi}{\partial s} \cdot \bar{\partial} \left(\frac{\partial \psi}{\partial t} \right) \wedge \partial \omega. \end{aligned}$$

In the same way, one deduces

$$d\Psi^2 = I^2 \cdot dt \wedge ds,$$

and

$$\frac{I^2}{b_2} = \int_X -\frac{\partial \psi}{\partial t} \cdot \partial \left(\frac{\partial \psi}{\partial s} \right) \wedge \bar{\partial} \omega + \int_X \frac{\partial \psi}{\partial s} \cdot \partial \left(\frac{\partial \psi}{\partial t} \right) \wedge \bar{\partial} \omega.$$

Combining above formulas, we have

$$(3.10) \quad -\frac{I^1}{a_2} + \frac{I^2}{b_2} = -\frac{I^0}{\sqrt{-1}}.$$

Setting $a_2 = -\sqrt{-1}$ and $b_2 = \sqrt{-1}$, we get

$$I^0 + I^1 + I^2 = 0.$$

Thus

$$(3.11) \quad d\Psi = 0$$

where

$$\Psi := \Psi^0 + \Psi^1 + \Psi^2.$$

The following theorem is an immediate consequence of the above discussion.

Theorem 3.1. *Let (X, g) be a compact complex surface and ω be its associated real $(1, 1)$ -form. The functional*

$$(3.12) \quad \begin{aligned} \mathcal{L}_\omega^M(\varphi', \varphi'') &:= \frac{1}{V_\omega} \int_0^1 \int_X \dot{\varphi}_t \cdot \omega_{\varphi_t}^2 dt \\ &\quad - \frac{1}{V_\omega} \int_0^1 \int_X \sqrt{-1} \partial \omega \wedge (\bar{\partial} \dot{\varphi}_t \cdot \varphi_t) dt \\ &\quad + \frac{1}{V_\omega} \int_0^1 \int_X \sqrt{-1} \bar{\partial} \omega \wedge (\partial \dot{\varphi}_t \cdot \varphi_t) dt \end{aligned}$$

is independent of the choice of the smooth path $\{\varphi_t\}_{0 \leq t \leq 1}$. In particular,

$$(3.13) \quad \begin{aligned} \mathcal{L}_\omega^M(\varphi) &:= \mathcal{L}_\omega(0, \varphi) = \frac{1}{3V_\omega} \int_X \varphi(\omega^2 + \omega \wedge \omega_\varphi + \omega_\varphi^2) \\ &+ \frac{1}{2V_\omega} \int_X \varphi[-\sqrt{-1}\partial\omega \wedge \bar{\partial}\varphi + \sqrt{-1}\bar{\partial}\omega \wedge \partial\varphi]. \end{aligned}$$

Proof. Applying Stocks' theorem to the region $\Delta = \{(s, t) \in \mathbb{R}^2 : 0 \leq s, t \leq 1\}$ and using equation (3.9), we have

$$\begin{aligned} 0 &= \int_\Delta d\Psi = \int_{\partial\Delta} \Psi = \int_{\partial\Delta} (\Psi^0 + \Psi^1 + \Psi^2) \\ &= \int_0^1 \dot{\varphi}_t \cdot \omega_{\varphi_t}^2 dt - \int_0^1 \int_X \frac{\partial\psi}{\partial s} \cdot \omega_\psi^2 ds \Big|_{t=0}^{t=1} \\ &+ \int_0^1 \int_X -\sqrt{-1}\partial\omega \wedge (\bar{\partial}\dot{\varphi}_t \cdot \varphi_t) dt \\ &- \int_0^1 \int_X -\sqrt{-1}\partial\omega \wedge \left(\bar{\partial} \left(\frac{\partial\psi}{\partial s} \right) \cdot \psi \right) ds \Big|_{t=0}^{t=1} \\ &+ \int_0^1 \int_X \sqrt{-1}\bar{\partial}\omega \wedge (\partial\dot{\varphi}_t \cdot \varphi_t) dt \\ &- \int_0^1 \int_X \sqrt{-1}\bar{\partial}\omega \wedge \left(\partial \left(\frac{\partial\psi}{\partial s} \right) \cdot \psi \right) ds \Big|_{t=0}^{t=1}. \end{aligned}$$

Equivalently,

$$\begin{aligned} \mathcal{L}_\omega^M(\varphi', \varphi'') &= \int_0^1 \int_X \frac{\partial\psi}{\partial s} \cdot \omega_\psi^2 ds \Big|_{t=0}^{t=1} \\ &+ \int_0^1 \int_X -\sqrt{-1}\partial\omega \wedge \left(\bar{\partial} \left(\frac{\partial\psi}{\partial s} \right) \cdot \psi \right) ds \Big|_{t=0}^{t=1} \\ &+ \int_0^1 \int_X \sqrt{-1}\bar{\partial}\omega \wedge \left(\partial \left(\frac{\partial\psi}{\partial s} \right) \cdot \psi \right) ds \Big|_{t=0}^{t=1}. \end{aligned}$$

It turns out that $\mathcal{L}_\omega^M(\varphi', \varphi'')$ is well-defined. For the second argument, we can choose the smooth path $\varphi_t = t \cdot \varphi$, $0 \leq t \leq 1$. $\square$

Remark 3.2. When (X, g) is a compact Kähler surface, the functional (3.12) or (3.13) coincides with the original one.

Definition 3.3. Suppose that S is a non-empty set and A is an additive group. A mapping $\mathcal{N} : S \times S \rightarrow A$ is said to satisfy the **1-cocycle condition** if

- (i) $\mathcal{N}(\sigma_1, \sigma_2) + \mathcal{N}(\sigma_2, \sigma_1) = 0$;
- (ii) $\mathcal{N}(\sigma_1, \sigma_2) + \mathcal{N}(\sigma_2, \sigma_3) + \mathcal{N}(\sigma_3, \sigma_1) = 0$.

Corollary 3.4. The functional $\mathcal{L}_\omega^M$ satisfies the 1-cocycle condition.

Corollary 3.5. For any $\varphi \in \mathcal{P}_\omega$ and any constant $C \in \mathbb{R}$, we have

$$(3.14) \quad \mathcal{L}_\omega^M(\varphi, \varphi + C) = C \cdot \left(1 - \frac{\text{Err}_\omega(\varphi)}{V_\omega} \right).$$

In particular, if $\partial\bar{\partial}\omega = 0$, then $\mathcal{L}_\omega^M(\varphi, \varphi + C) = C$.

Proof. We choose the smooth path $\varphi_t = \varphi + t \cdot C$, $t \in [0, 1]$. So

$$\begin{aligned} \mathcal{L}_\omega^M(\varphi, \varphi + C) &= \frac{1}{V_\omega} \int_0^1 \int_X C \cdot \omega_{\varphi+C \cdot t}^2 dt = \frac{1}{V_\omega} \int_0^1 \int_X C \cdot \omega_\varphi^2 dt \\ &= \frac{1}{V_\omega} \int_X C \cdot \omega_\varphi^2 = \frac{C}{V_\omega} \cdot V_\omega(\varphi) = C \cdot \left(1 - \frac{\text{Err}_\omega(\varphi)}{V_\omega}\right). \end{aligned}$$

If furthermore $\partial\bar{\partial}\omega = 0$, then $\text{Err}_\omega(\varphi) = 0$ for any $\varphi \in \mathcal{P}_\omega$. $\square$

Corollary 3.6. *For any $\varphi_1, \varphi_2 \in \mathcal{P}_\omega$ and any constant $C \in \mathbb{R}$, we have*

$$(3.15) \quad \mathcal{L}_\omega^M(\varphi_1, \varphi_2 + C) = \mathcal{L}_\omega^M(\varphi_1, \varphi_2) + C \cdot \left(1 - \frac{\text{Err}_\omega(\varphi_2)}{V_\omega}\right).$$

Proof. From Corollary 3.4, one has

$$\mathcal{L}_\omega^M(\varphi_1, \varphi_2 + C) + \mathcal{L}_\omega^M(\varphi_2, \varphi_1) = \mathcal{L}_\omega^M(\varphi_2, \varphi_2 + C).$$

Then the conclusion follows from Corollary 3.5. $\square$

4. AUBIN-YAU FUNCTIONALS ON COMPACT COMPLEX SURFACES

In this section we extend Aubin-Yau functionals to compact complex surfaces, including Kähler surfaces, and deduce a number of basic properties of these functionals.

4.1. Aubin-Yau functionals on compact Kähler manifolds. Suppose that (X, ω) is a compact Kähler manifold of dimension n . For $(\varphi', \varphi'') \in \mathcal{P}_\omega \times \mathcal{P}_\omega$, Aubin-Yau functionals are defined by

$$(4.1) \quad \mathcal{I}_\omega^{\text{AY}}(\varphi', \varphi'') = \frac{1}{V_\omega} \int_X (\varphi'' - \varphi')(\omega_{\varphi'}^n - \omega_{\varphi''}^n),$$

$$(4.2) \quad \mathcal{J}_\omega^{\text{AY}}(\varphi', \varphi'') = -\mathcal{L}_\omega^M(\varphi', \varphi'') + \frac{1}{V_\omega} \int_X (\varphi'' - \varphi')\omega_{\varphi'}^n.$$

By definition, we have

$$(4.3) \quad \mathcal{J}_\omega^{\text{AY}}(\varphi', \varphi'') + \mathcal{J}_\omega^{\text{AY}}(\varphi'', \varphi') = \mathcal{I}_\omega^{\text{AY}}(\varphi', \varphi'') = \mathcal{I}_\omega^{\text{AY}}(\varphi'', \varphi').$$

For any $\varphi \in \mathcal{P}_\omega$ we set

$$(4.4) \quad \mathcal{I}_\omega^{\text{AY}}(\varphi) = \frac{1}{V_\omega} \int_X \varphi(\omega^n - \omega_\varphi^n),$$

$$(4.5) \quad \mathcal{J}_\omega^{\text{AY}}(\varphi) = \int_0^1 \frac{\mathcal{I}_\omega(s \cdot \varphi)}{s} ds = \frac{1}{V_\omega} \int_0^1 \int_X \varphi(\omega^n - \omega_{s \cdot \varphi}^n) ds.$$

It's clear that

$$(4.6) \quad \mathcal{I}_\omega^{\text{AY}}(\varphi) = \mathcal{I}_\omega^{\text{AY}}(0, \varphi), \quad \mathcal{J}_\omega^{\text{AY}}(\varphi) = \mathcal{J}_\omega^{\text{AY}}(0, \varphi).$$

By definition we have

$$\begin{aligned}
& \mathcal{J}_\omega^{\text{AY}}(\varphi) \\
&= \frac{1}{V_\omega} \int_0^1 ds \int_X \varphi(-\sqrt{-1} \partial \bar{\partial}(s \cdot \varphi)) \wedge \sum_{i=1}^{n-1} \omega^{n-1-i} \wedge \omega_{s \cdot \varphi}^i \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_0^1 s \cdot ds \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{i=0}^{n-1} \omega^{n-1-i} \wedge [\omega + s(\omega_\varphi - \omega)]^i \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_0^1 s \cdot ds \int_X \varphi \cdot \partial \bar{\partial} \varphi \sum_{i=0}^{n-1} \omega^{n-1-i} \wedge \sum_{j=0}^i \binom{i}{j} (1-s)^{i-j} s^j \omega^{i-j} \wedge \omega_\varphi^j \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{j=0}^{n-1} \sum_{i=j}^{n-1} \binom{i}{j} \omega^{n-1-j} \wedge \omega_\varphi^j \int_0^1 (1-s)^{i-j} s^{1+j} \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{j=0}^{n-1} \omega^{n-1-j} \wedge \omega_\varphi^j \cdot \sum_{i=j}^{n-1} \binom{i}{j} \cdot \frac{(i-j)! \cdot (j+1)!}{(i+2)!} \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{j=0}^{n-1} \omega^{n-1-j} \wedge \omega_\varphi^j \cdot \sum_{i=j}^{n-1} \frac{j+1}{(i+2)(i+1)} \\
&= \frac{-\sqrt{-1}}{V_\omega} \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{j=0}^{n-1} \frac{n-j}{n+1} \omega^{n-1-j} \wedge \omega_\varphi^j
\end{aligned}$$

since

$$\sum_{i=j}^{n-1} \frac{1}{(i+2)(i+1)} = \sum_{i=j}^{n-1} \left(\frac{1}{i+1} - \frac{1}{i+2} \right) = \frac{1}{j+1} - \frac{1}{n+1}.$$

On the other hand,

$$\frac{n}{n+1} \mathcal{I}_\omega^{\text{AY}}(\varphi) = \frac{-\sqrt{-1}}{V_\omega} \int_X \varphi \cdot \partial \bar{\partial} \varphi \wedge \sum_{i=0}^{n-1} \frac{n}{n+1} \omega^{n-1-i} \wedge \omega_\varphi^i$$

Hence

$$\begin{aligned}
(4.7) \quad & \frac{n}{n+1} \mathcal{I}_\omega^{\text{AY}}(\varphi) - \mathcal{J}_\omega^{\text{AY}}(\varphi) \\
&= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \sum_{j=1}^{n-1} \frac{j}{n+1} \omega^{n-1-j} \wedge \omega_\varphi^j.
\end{aligned}$$

Moreover,

$$\begin{aligned}
(4.8) \quad & (n+1) \mathcal{J}_\omega^{\text{AY}}(\varphi) - \mathcal{I}_\omega^{\text{AY}}(\varphi) \\
&= \frac{1}{V_\omega} \int_X \varphi \cdot -\sqrt{-1} \partial \bar{\partial} \varphi \wedge \sum_{j=0}^{n-1} (n-1-j) \omega^{n-1-j} \wedge \omega_\varphi^j.
\end{aligned}$$

Remark 4.1. Notice that formulae (4.7) and (4.8) are also valid when ω is non-Kähler.

4.2. Aubin-Yau functionals over compact complex surfaces. Let (X, g) be a compact complex manifold of the complex dimension n and ω be its associated real $(1, 1)$ -form. From Remark 4.1, we can formally use the notion $\mathcal{I}_\omega^{\text{AY}}$, $\mathcal{J}_\omega^{\text{AY}}$, and $\mathcal{L}_\omega^{\text{M}}$, but now ω may not be Kähler. Precisely, for any $\varphi \in \mathcal{P}_\omega$ we set

$$(4.9) \quad \mathcal{I}_{\omega|\bullet}^{\text{AY}}(\varphi) = \frac{1}{V_\omega} \int_X \varphi(\omega^n - \omega_\varphi^n),$$

$$(4.10) \quad \mathcal{J}_{\omega|\bullet}^{\text{AY}}(\varphi) = \int_0^1 \frac{\mathcal{I}_{\omega|\bullet}^{\text{AY}}(s \cdot \varphi)}{s} ds = \frac{1}{V_\omega} \int_0^1 \int_X \varphi(\omega^n - \omega_{s \cdot \varphi}^n) ds.$$

Hence

$$(4.11) \quad \begin{aligned} & \frac{n}{n+1} \mathcal{I}_{\omega|\bullet}^{\text{AY}}(\varphi) - \mathcal{J}_{\omega|\bullet}^{\text{AY}}(\varphi) \\ &= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \sum_{j=1}^{n-1} \frac{j}{n+1} \omega^{n-1-j} \wedge \omega_\varphi^j. \end{aligned}$$

Moreover,

$$(4.12) \quad \begin{aligned} & (n+1) \mathcal{J}_{\omega|\bullet}^{\text{AY}}(\varphi) - \mathcal{I}_{\omega|\bullet}^{\text{AY}}(\varphi) \\ &= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \sum_{j=0}^{n-1} (n-1-j) \omega^{n-1-j} \wedge \omega_\varphi^j. \end{aligned}$$

Restricting to compact complex surfaces and introducing two extra functionals on $\mathcal{P}_\omega$

$$(4.13) \quad \mathcal{A}_\omega(\varphi) := \frac{1}{2V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \omega \wedge \bar{\partial} \varphi),$$

$$(4.14) \quad \mathcal{B}_\omega(\varphi) := \frac{1}{2V_\omega} \int_X \varphi \cdot \sqrt{-1} \partial \omega \wedge \bar{\partial} \varphi,$$

(clearly $\overline{\mathcal{A}_\omega(\varphi)} = \mathcal{B}_\omega(\varphi)$), we define Aubin-Yau functionals as follows (Here constants a, b, c, d are determined later, and actually $a = b = c = d = 2$)

$$(4.15) \quad \mathcal{I}_\omega^{\text{AY}}(\varphi) := \mathcal{I}_{\omega|\bullet}^{\text{AY}}(\varphi) + a \mathcal{A}_\omega(\varphi) + b \mathcal{B}_\omega(\varphi),$$

$$(4.16) \quad \mathcal{J}_\omega^{\text{AY}}(\varphi) := -\mathcal{L}_\omega^{\text{M}}(\varphi) + \frac{1}{V_\omega} \int_X \varphi \cdot \omega^2 + c \mathcal{A}_\omega(\varphi) + d \mathcal{B}_\omega(\varphi).$$

Since

$$\begin{aligned} \mathcal{J}_{\omega|\bullet}^{\text{AY}}(\varphi) &= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \sum_{j=0}^1 \frac{2-j}{3} (\omega^{1-j} \wedge \omega_\varphi^j) \\ &= \frac{1}{V_\omega} \int_X \varphi (\omega - \omega_\varphi) \wedge \left(\frac{2}{3} \omega + \frac{1}{3} \omega_\varphi \right) \\ &= \frac{1}{V_\omega} \int_X \varphi \left(\frac{2}{3} \omega^2 - \frac{1}{3} \omega \wedge \omega_\varphi - \frac{1}{3} \omega_\varphi^2 \right), \end{aligned}$$

it follows that

$$(4.17) \quad \mathcal{J}_\omega^{\text{AY}}(\varphi) = \mathcal{J}_{\omega|\bullet}^{\text{AY}}(\varphi) + (c-1) \mathcal{A}_\omega(\varphi) + (d-1) \mathcal{B}_\omega(\varphi).$$

Using Remark 4.1 or the previous subsection, we deduce

$$\begin{aligned}
& \frac{2}{3}(\mathcal{I}_\omega^{\text{AY}}(\varphi) - a\mathcal{A}_\omega(\varphi) - b\mathcal{B}_\omega(\varphi)) - (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1)\mathcal{A}_\omega(\varphi) - (d-1)\mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{V_\omega} \int_X \varphi(-\sqrt{-1}\partial\bar{\partial}\varphi) \sum_{j=1}^1 \frac{j}{3} \omega^{1-j} \wedge \omega_\varphi^j \\
&= \frac{1}{V_\omega} \int_X \varphi \cdot \frac{1}{3} \omega_\varphi \wedge (-\sqrt{-1}\partial\bar{\partial}\varphi) = \frac{1}{V_\omega} \int_X \varphi \cdot \frac{1}{3} \omega_\varphi \wedge \sqrt{-1}\partial\bar{\partial}\varphi.
\end{aligned}$$

Thus the left hand side has two slightly different expressions. If we adopt the first one, we have

$$\begin{aligned}
& \frac{2}{3}(\mathcal{I}_\omega^{\text{AY}}(\varphi) - a\mathcal{A}_\omega(\varphi) - b\mathcal{B}_\omega(\varphi)) - (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1)\mathcal{A}_\omega(\varphi) - (d-1)\mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{3V_\omega} \int_X \sqrt{-1}\partial(\varphi \cdot \omega_\varphi) \wedge \bar{\partial}\varphi = \frac{1}{3V_\omega} \int_X \sqrt{-1}(\partial\varphi \wedge \omega_\varphi + \varphi \cdot \partial\omega) \wedge \bar{\partial}\varphi \\
&= \frac{1}{3V_\omega} \int_X \sqrt{-1}\partial\varphi \wedge \bar{\partial}\varphi \wedge \omega_\varphi - \frac{2}{3}\mathcal{A}_\omega(\varphi).
\end{aligned}$$

On the other hand, using the second expression gives

$$\begin{aligned}
& \frac{2}{3}(\mathcal{I}_\omega^{\text{AY}}(\varphi) - a\mathcal{A}_\omega(\varphi) - b\mathcal{B}_\omega(\varphi)) - (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1)\mathcal{A}_\omega(\varphi) - (d-1)\mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{3V_\omega} \int_X -\sqrt{-1}\bar{\partial}(\varphi \cdot \omega_\varphi) \wedge \partial\varphi = \frac{1}{3V_\omega} \int_X -\sqrt{-1}(\bar{\partial}\varphi \wedge \omega_\varphi + \varphi \cdot \bar{\partial}\omega) \wedge \partial\varphi \\
&= \frac{1}{3V_\omega} \int_X \sqrt{-1}\partial\varphi \wedge \bar{\partial}\varphi \wedge \omega_\varphi - \frac{2}{3}\mathcal{B}_\omega(\varphi).
\end{aligned}$$

Therefore

$$\begin{aligned}
& \frac{2}{3}(\mathcal{I}_\omega^{\text{AY}}(\varphi) - a\mathcal{A}_\omega(\varphi) - b\mathcal{B}_\omega(\varphi)) - (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1)\mathcal{A}_\omega(\varphi) - (d-1)\mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{3V_\omega} \int_X \sqrt{-1}\partial\varphi \wedge \bar{\partial}\varphi \wedge \omega_\varphi - \frac{\mathcal{A}_\omega(\varphi) + \mathcal{B}_\omega(\varphi)}{3},
\end{aligned}$$

or, equivalently,

$$(4.18) \quad \frac{2}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi) - \mathcal{J}_\omega^{\text{AY}}(\varphi) = \frac{1}{3V_\omega} \int_X \sqrt{-1}\partial\varphi \wedge \bar{\partial}\varphi \wedge \omega_\varphi$$

where we require

$$(4.19) \quad \frac{2}{3}a - (c-1) - \frac{1}{3} = 0 = \frac{2}{3}b - (d-1) - \frac{1}{3}.$$

Theorem 4.2. *For any $\varphi \in \mathcal{P}_\omega$, one has*

$$(4.20) \quad \frac{2}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi) - \mathcal{J}_\omega^{\text{AY}}(\varphi) \geq 0.$$

Using (4.12) yields

$$\begin{aligned}
& 3 \left(\mathcal{J}_\omega^{\text{AY}}(\varphi) - \frac{1}{2} \mathcal{A}_\omega(\varphi) - \frac{1}{2} \mathcal{B}_\omega(\varphi) \right) - (\mathcal{I}_\omega^{\text{AY}}(\varphi) - \mathcal{A}_\omega(\varphi) - \mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \sum_{j=0}^1 (1-j) \omega^{1-j} \wedge \omega_\varphi^j \\
&= \frac{1}{V_\omega} \int_X \varphi \cdot (-\sqrt{-1} \partial \bar{\partial} \varphi) \wedge \omega \\
&= \frac{1}{V_\omega} \int_X (\varphi \cdot \omega) \wedge (-\sqrt{-1} \partial \bar{\partial} \varphi) = \frac{1}{V_\omega} \int_X (\varphi \cdot \omega) \wedge \sqrt{-1} \partial \bar{\partial} \varphi.
\end{aligned}$$

As the proof of Theorem 4.2, we have

$$\begin{aligned}
& 3 (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1) \mathcal{A}_\omega(\varphi) - (d-1) \mathcal{B}_\omega(\varphi)) - (\mathcal{I}_\omega^{\text{AY}}(\varphi) - a \mathcal{A}_\omega(\varphi) - b \mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{V_\omega} \int_X \sqrt{-1} \partial (\varphi \cdot \omega) \wedge \bar{\partial} \varphi = \frac{1}{V_\omega} \int_X \sqrt{-1} (\partial \varphi \wedge \omega + \varphi \cdot \partial \omega) \wedge \bar{\partial} \varphi \\
&= \frac{1}{V_\omega} \int_X \sqrt{-1} \partial \varphi \wedge \bar{\partial} \varphi \wedge \omega - 2 \mathcal{A}_\omega(\varphi)
\end{aligned}$$

and

$$\begin{aligned}
& 3 (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1) \mathcal{A}_\omega(\varphi) - (d-1) \mathcal{B}_\omega(\varphi)) - (\mathcal{I}_\omega^{\text{AY}}(\varphi) - a \mathcal{A}_\omega(\varphi) - b \mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{V_\omega} \int_X -\sqrt{-1} \bar{\partial} (\varphi \cdot \omega) \wedge \partial \varphi = \frac{1}{V_\omega} \int_X -\sqrt{-1} (\bar{\partial} \varphi \wedge \omega + \varphi \cdot \bar{\partial} \omega) \wedge \partial \varphi \\
&= \frac{1}{V_\omega} \int_X \sqrt{-1} \bar{\partial} \varphi \wedge \partial \varphi \wedge \omega - 2 \mathcal{B}_\omega(\varphi).
\end{aligned}$$

Hence

$$\begin{aligned}
& 3 (\mathcal{J}_\omega^{\text{AY}}(\varphi) - (c-1) \mathcal{A}_\omega(\varphi) - (d-1) \mathcal{B}_\omega(\varphi)) - (\mathcal{I}_\omega^{\text{AY}}(\varphi) - a \mathcal{A}_\omega(\varphi) - b \mathcal{B}_\omega(\varphi)) \\
&= \frac{1}{V_\omega} \int_X \sqrt{-1} \partial \varphi \wedge \bar{\partial} \varphi \wedge \omega - (\mathcal{A}_\omega(\varphi) + \mathcal{B}_\omega(\varphi)).
\end{aligned}$$

Equivalently,

$$(4.21) \quad 3 \mathcal{J}_\omega^{\text{AY}}(\varphi) - \mathcal{I}_\omega^{\text{AY}}(\varphi) = \frac{1}{V_\omega} \int_X \sqrt{-1} \partial \varphi \wedge \bar{\partial} \varphi \wedge \omega,$$

where we also require

$$(4.22) \quad 3(c-1) - a - 1 = 0 = 3(d-1) - b - 1.$$

Theorem 4.3. *For any $\varphi \in \mathcal{P}_\omega$, one has*

$$(4.23) \quad 3 \mathcal{J}_\omega^{\text{AY}}(\varphi) - \mathcal{I}_\omega^{\text{AY}}(\varphi) \geq 0.$$

Combining (4.19) and (4.22) we obtain the value of those constants:

$$a = b = c = d = 2.$$

Corollary 4.4. *For any compact complex surface (X, g) and any real-valued smooth function $\varphi \in \mathcal{P}_\omega$, we have*

$$\begin{aligned} \frac{1}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi) &\leq \mathcal{J}_\omega^{\text{AY}}(\varphi) \leq \frac{2}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi), \\ \frac{3}{2}\mathcal{J}_\omega^{\text{AY}}(\varphi) &\leq \mathcal{I}_\omega(\varphi) \leq 3\mathcal{J}_\omega^{\text{AY}}(\varphi), \\ \frac{1}{2}\mathcal{J}_\omega^{\text{AY}}(\varphi) &\leq \frac{1}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi) \leq \mathcal{I}_\omega^{\text{AY}}(\varphi) - \mathcal{J}_\omega^{\text{AY}}(\varphi) \leq \frac{2}{3}\mathcal{I}_\omega^{\text{AY}}(\varphi) \leq 2\mathcal{J}_\omega^{\text{AY}}(\varphi), \end{aligned}$$

where ω is its associated real $(1, 1)$ -form.

REFERENCES

1. Chen, X. X., Tian, G., *Ricci flow on Kähler-Einstein surfaces*, Invent. Math. **147**(2002), no.3, 487–544.
2. Dinew, S., Kolodziej, S., *Pluripotential estimates on compact Hermitian manifolds*, preprint, arXiv: 0910.3937, 2009.
3. Gauduchon, P., *Sur la 1-forme de torsion d'une variété hermitienne compacte*, Math. Ann. **267** (1984), 495–518.
4. Guan, B., Li, Q., *Complex Monge-Ampère equations on Hermitian manifold*, preprint, arXiv: 0906.3548, 2009.
5. Lübke, M., Teleman, A., *The Kobayashi-Hitchin correspondence*, World Scientific, 1995.
6. Mabuchi, T., *K-energy maps integrating Futaki invariants*, Tohoku Math. Journ., **38**(1986), 575–593.
7. Phong, D.H., Sturm, J., *Lectures on stability and constant scalar curvature*, preprint, arXiv: 0801.4179, 2008.
8. Siu, Y.T., *Every K3-surface is Kähler*, Invent. Math., **73** (1983), 139–150.
9. Tosatti, V., Weinkove, B., *Estimates for the complex Monge-Ampère equation on Hermitian and balanced manifolds*, preprint, arXiv: 0909.4496, 2009.
10. Tosatti, V., Weinkove, B., *The complex Monge-ampère equation on compact Hermitian manifolds*, preprint, arXiv: 0910.1390, 2009.
11. Yau, S.-T., *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I*, Comm. Pure. Appl. Math., **31** (1978), no.3, 339–411.
12. Yau, S.-T., *Review of geometry and analysis*, Kodaira's issue. Asian J. Math. **4**(2000), no.1, 235–278

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA 02138
E-mail address: yili@math.harvard.edu